

**CDA Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP2868b****Specification**

---

**CDA Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [P32320](#)**CDA Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 978**Other Names**

Cytidine deaminase, Cytidine aminohydrolase, CDA, CDD

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP2868b](/products/AP2868b) was selected from the C-term region of human CDA. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**CDA Antibody (C-term) Blocking Peptide - Protein Information****Name** CDA ([HGNC:1712](#))**Synonyms** CDD**Function**

This enzyme scavenges exogenous and endogenous cytidine and 2'-deoxycytidine for UMP synthesis.

**Tissue Location**

Highly expressed in granulocytes while expression is very low in fibroblasts, chondrocytes, monocytes, and T- as well as B-cell lines

**CDA Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **CDA Antibody (C-term) Blocking Peptide - Images**

#### **CDA Antibody (C-term) Blocking Peptide - Background**

CDA forms a homotetramer that catalyzes the irreversible hydrolytic deamination of cytidine and deoxycytidine to uridine and deoxyuridine, respectively. This protein is one of several deaminases responsible for maintaining the cellular pyrimidine pool. Mutations in the CDA gene are associated with decreased sensitivity to the cytosine nucleoside analogue cytosine arabinoside used in the treatment of certain childhood leukemias.

#### **CDA Antibody (C-term) Blocking Peptide - References**

Laliberte J., Momparler R.L. Cancer Res. 54:5401-5407(1994) Demontis S., Terao M. Biochim. Biophys. Acta 1443:323-333(1998) Kuhn K., Bertling W.M., Emmrich F. Biochem. Biophys. Res. Commun. 190:1-7(1993)