

**Phospho-Aura-T288 Antibody Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP3030a****Specification**

---

**Phospho-Aura-T288 Antibody Blocking Peptide - Product Information**

Primary Accession [O14965](#)  
Other Accession [NP\\_003591](#)

**Phospho-Aura-T288 Antibody Blocking Peptide - Additional Information**

**Gene ID** 6790

**Other Names**

Aurora kinase A, Aurora 2, Aurora/IPL1-related kinase 1, ARK-1, Aurora-related kinase 1, hARK1, Breast tumor-amplified kinase, Serine/threonine-protein kinase 15, Serine/threonine-protein kinase 6, Serine/threonine-protein kinase aurora-A, AURKA

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP3030a](/product/products/AP3030a) was selected from the region encoding phosphorylated tyrosine at position 288 of human Aura. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Phospho-Aura-T288 Antibody Blocking Peptide - Protein Information**

**Name** AURKA ([HGNC:11393](#))

**Function**

Mitotic serine/threonine kinase that contributes to the regulation of cell cycle progression (PubMed: [26246606](http://www.uniprot.org/citations/26246606), PubMed: [12390251](http://www.uniprot.org/citations/12390251), PubMed: [18615013](http://www.uniprot.org/citations/18615013), PubMed: [11039908](http://www.uniprot.org/citations/11039908), PubMed: [17125279](http://www.uniprot.org/citations/17125279), PubMed: [17360485](http://www.uniprot.org/citations/17360485)). Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome

duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis (PubMed:<a href="http://www.uniprot.org/citations/26246606" target="\_blank">26246606</a>, PubMed:<a href="http://www.uniprot.org/citations/14523000" target="\_blank">14523000</a>). Required for normal spindle positioning during mitosis and for the localization of NUMA1 and DCTN1 to the cell cortex during metaphase (PubMed:<a href="http://www.uniprot.org/citations/27335426" target="\_blank">27335426</a>). Required for initial activation of CDK1 at centrosomes (PubMed:<a href="http://www.uniprot.org/citations/13678582" target="\_blank">13678582</a>, PubMed:<a href="http://www.uniprot.org/citations/15128871" target="\_blank">15128871</a>). Phosphorylates numerous target proteins, including ARHGEF2, BORA, BRCA1, CDC25B, DLGP5, HDAC6, KIF2A, LATS2, NDEL1, PARD3, PPP1R2, PLK1, RASSF1, TACC3, p53/TP53 and TPX2 (PubMed:<a href="http://www.uniprot.org/citations/18056443" target="\_blank">18056443</a>, PubMed:<a href="http://www.uniprot.org/citations/15128871" target="\_blank">15128871</a>, PubMed:<a href="http://www.uniprot.org/citations/14702041" target="\_blank">14702041</a>, PubMed:<a href="http://www.uniprot.org/citations/11551964" target="\_blank">11551964</a>, PubMed:<a href="http://www.uniprot.org/citations/15147269" target="\_blank">15147269</a>, PubMed:<a href="http://www.uniprot.org/citations/15987997" target="\_blank">15987997</a>, PubMed:<a href="http://www.uniprot.org/citations/17604723" target="\_blank">17604723</a>, PubMed:<a href="http://www.uniprot.org/citations/18615013" target="\_blank">18615013</a>). Regulates KIF2A tubulin depolymerase activity (PubMed:<a href="http://www.uniprot.org/citations/19351716" target="\_blank">19351716</a>). Important for microtubule formation and/or stabilization (PubMed:<a href="http://www.uniprot.org/citations/18056443" target="\_blank">18056443</a>). Required for normal axon formation (PubMed:<a href="http://www.uniprot.org/citations/19812038" target="\_blank">19812038</a>). Plays a role in microtubule remodeling during neurite extension (PubMed:<a href="http://www.uniprot.org/citations/19668197" target="\_blank">19668197</a>). Also acts as a key regulatory component of the p53/TP53 pathway, and particularly the checkpoint- response pathways critical for oncogenic transformation of cells, by phosphorylating and destabilizing p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/14702041" target="\_blank">14702041</a>). Phosphorylates its own inhibitors, the protein phosphatase type 1 (PP1) isoforms, to inhibit their activity (PubMed:<a href="http://www.uniprot.org/citations/11551964" target="\_blank">11551964</a>). Inhibits cilia outgrowth (By similarity). Required for cilia disassembly via phosphorylation of HDAC6 and subsequent deacetylation of alpha-tubulin (PubMed:<a href="http://www.uniprot.org/citations/17604723" target="\_blank">17604723</a>, PubMed:<a href="http://www.uniprot.org/citations/20643351" target="\_blank">20643351</a>). Regulates protein levels of the anti-apoptosis protein BIRC5 by suppressing the expression of the SCF(FBXL7) E3 ubiquitin-protein ligase substrate adapter FBXL7 through the phosphorylation of the transcription factor FOXP1 (PubMed:<a href="http://www.uniprot.org/citations/28218735" target="\_blank">28218735</a>).

### Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:P97477}. Cell projection, neuron projection {ECO:0000250|UniProtKB:P97477}. Cell projection, cilium. Cytoplasm, cytoskeleton, cilium basal body. Basolateral cell membrane {ECO:0000250|UniProtKB:F1PNY0}. Note=Detected at the neurite hillock in developing neurons (By similarity). Localizes at the centrosome in mitotic cells from early prophase until telophase, but also localizes to the spindle pole MTs from prophase to anaphase (PubMed:9606188, PubMed:17229885, PubMed:21225229). Colocalized with SIRT2 at centrosome (PubMed:22014574). Moves to the midbody during both telophase and cytokinesis (PubMed:17726514). Associates with both the pericentriolar material (PCM) and centrioles (PubMed:22014574). The localization to the spindle poles is regulated by AAAS (PubMed:26246606) {ECO:0000250|UniProtKB:P97477, ECO:0000269|PubMed:17229885, ECO:0000269|PubMed:17726514, ECO:0000269|PubMed:21225229, ECO:0000269|PubMed:22014574, ECO:0000269|PubMed:26246606, ECO:0000269|PubMed:9606188}

**Tissue Location**

Highly expressed in testis and weakly in skeletal muscle, thymus and spleen. Also highly expressed in colon, ovarian, prostate, neuroblastoma, breast and cervical cancer cell lines

**Phospho-Aura-T288 Antibody Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**Phospho-Aura-T288 Antibody Blocking Peptide - Images****Phospho-Aura-T288 Antibody Blocking Peptide - Background**

The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript variants encoding the same protein have been found for this gene.

**Phospho-Aura-T288 Antibody Blocking Peptide - References**

Kamada, K., et al., Oncol. Rep. 12(3):593-599 (2004).Satinover, D.L., et al., Proc. Natl. Acad. Sci. U.S.A. 101(23):8625-8630 (2004).Ouchi, M., et al., J. Biol. Chem. 279(19):19643-19648 (2004).Kunitoku, N., et al., Dev. Cell 5(6):853-864 (2003).Prigent, C., et al., Cell 114(5):531-532 (2003).