

Phospho-CDC6(S54) Antibody Blocking peptide
Synthetic peptide
Catalog # BP3058a**Specification**

Phospho-CDC6(S54) Antibody Blocking peptide - Product InformationPrimary Accession [Q99741](#)**Phospho-CDC6(S54) Antibody Blocking peptide - Additional Information****Gene ID** 990**Other Names**

Cell division control protein 6 homolog, CDC6-related protein, Cdc18-related protein, HsCdc18, p62(cdc6), HsCDC6, CDC6, CDC18L

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP3058a](/product/products/AP3058a) was selected from the 47-61 region of human Phospho-CDC6-S54. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-CDC6(S54) Antibody Blocking peptide - Protein Information**Name** CDC6**Synonyms** CDC18L**Function**

Involved in the initiation of DNA replication. Also participates in checkpoint controls that ensure DNA replication is completed before mitosis is initiated.

Cellular Location

Nucleus. Cytoplasm Note=The protein is nuclear in G1 and cytoplasmic in S-phase cells (PubMed:9566895).

Phospho-CDC6(S54) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Phospho-CDC6(S54) Antibody Blocking peptide - Images

Phospho-CDC6(S54) Antibody Blocking peptide - Background

The protein encoded by this gene is highly similar to *Saccharomyces cerevisiae* Cdc6, a protein essential for the initiation of DNA replication. This protein functions as a regulator at the early steps of DNA replication. It localizes in cell nucleus during cell cycle G1, but translocates to the cytoplasm at the start of S phase. The subcellular translocation of this protein during cell cycle is regulated through its phosphorylation by Cdks. Transcription of this protein was reported to be regulated in response to mitogenic signals through transcriptional control mechanism involving E2F proteins.

Phospho-CDC6(S54) Antibody Blocking peptide - References

Alexandrow, M.G., et al., Mol. Cell. Biol. 24(4):1614-1627 (2004). Yim, H., et al., Mol. Biol. Cell 14(10):4250-4259 (2003). Clay-Farrace, L., et al., EMBO J. 22(3):704-712 (2003). Pelizon, C., et al., EMBO Rep. 3(8):780-784 (2002). Bermejo, R., et al., Mol. Biol. Cell 13(11):3989-4000 (2002).