

Phospho-ErbB2(Y1248) Antibody Blocking peptide
Synthetic peptide
Catalog # BP3094a**Specification**

Phospho-ErbB2(Y1248) Antibody Blocking peptide - Product InformationPrimary Accession [P04626](#)**Phospho-ErbB2(Y1248) Antibody Blocking peptide - Additional Information****Gene ID** 2064**Other Names**

Receptor tyrosine-protein kinase erbB-2, Metastatic lymph node gene 19 protein, MLN 19, Proto-oncogene Neu, Proto-oncogene c-ErbB-2, Tyrosine kinase-type cell surface receptor HER2, p185erbB2, CD340, ERBB2, HER2, MLN19, NEU, NGL

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP3094a](/product/products/AP3094a) was selected from the region of human Phospho-ErbB2-Y1248. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-ErbB2(Y1248) Antibody Blocking peptide - Protein Information**Name** ERBB2**Synonyms** HER2, MLN19, NEU, NGL**Function**

Protein tyrosine kinase that is part of several cell surface receptor complexes, but that apparently needs a coreceptor for ligand binding. Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Regulates outgrowth and stabilization of peripheral microtubules (MTs). Upon ERBB2 activation, the MEMO1-RHOA-DIAPH1 signaling pathway elicits the phosphorylation and thus the inhibition of GSK3B at cell membrane. This prevents the phosphorylation of APC and CLASP2, allowing its association with the cell membrane. In turn, membrane-bound APC allows the localization of MACF1 to the cell membrane, which is required for microtubule capture and stabilization.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell projection, ruffle membrane; Single-pass type I membrane protein. Note=Internalized from the cell membrane in response to EGF stimulation. [Isoform 2]: Cytoplasm. Nucleus.

Tissue Location

Expressed in a variety of tumor tissues including primary breast tumors and tumors from small bowel, esophagus, kidney and mouth.

Phospho-ErbB2(Y1248) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Phospho-ErbB2(Y1248) Antibody Blocking peptide - Images**Phospho-ErbB2(Y1248) Antibody Blocking peptide - Background**

ErbB2 is a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors.

Phospho-ErbB2(Y1248) Antibody Blocking peptide - References

Provinciali, M., et al., Int. J. Cancer 115(1):36-45 (2005). Yee, L.D., et al., J. Nutr. 135(5):983-988 (2005). Beckers, J., et al., Int. J. Cancer 114(4):590-597 (2005). Brandt, B., et al., Biochem. Biophys. Res. Commun. 329(1):318-323 (2005). Ghatak, S., et al., J. Biol. Chem. 280(10):8875-8883 (2005).