

**Phospho-CHUK(T23) Antibody Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP3374a****Specification**

---

**Phospho-CHUK(T23) Antibody Blocking peptide - Product Information**Primary Accession [O15111](#)**Phospho-CHUK(T23) Antibody Blocking peptide - Additional Information****Gene ID** 1147**Other Names**

Inhibitor of nuclear factor kappa-B kinase subunit alpha, I-kappa-B kinase alpha, IKK-A, IKK-alpha, IKBKA, IkappaB kinase, Conserved helix-loop-helix ubiquitous kinase, I-kappa-B kinase 1, IKK1, Nuclear factor NF-kappa-B inhibitor kinase alpha, NFKBKA, Transcription factor 16, TCF-16, CHUK, IKKA, TCF16

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP3374a](/product/products/AP3374a) was selected from the region of human Phospho-CHUK-T23. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Phospho-CHUK(T23) Antibody Blocking peptide - Protein Information****Name** CHUK**Synonyms** IKKA, TCF16**Function**

Serine kinase that plays an essential role in the NF-kappa-B signaling pathway which is activated by multiple stimuli such as inflammatory cytokines, bacterial or viral products, DNA damages or other cellular stresses (PubMed: [9244310](http://www.uniprot.org/citations/9244310), PubMed: [9252186](http://www.uniprot.org/citations/9252186), PubMed: [9346484](http://www.uniprot.org/citations/9346484), PubMed: [18626576](http://www.uniprot.org/citations/18626576)). Acts as a part of the canonical IKK complex in the conventional

pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B on serine residues (PubMed:<a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed:<a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed:<a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>, PubMed:<a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>, PubMed:<a href="http://www.uniprot.org/citations/35952808" target="\_blank">35952808</a>). These modifications allow polyubiquitination of the inhibitors and subsequent degradation by the proteasome (PubMed:<a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed:<a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed:<a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>, PubMed:<a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>). In turn, free NF-kappa-B is translocated into the nucleus and activates the transcription of hundreds of genes involved in immune response, growth control, or protection against apoptosis (PubMed:<a href="http://www.uniprot.org/citations/9244310" target="\_blank">9244310</a>, PubMed:<a href="http://www.uniprot.org/citations/9252186" target="\_blank">9252186</a>, PubMed:<a href="http://www.uniprot.org/citations/9346484" target="\_blank">9346484</a>, PubMed:<a href="http://www.uniprot.org/citations/18626576" target="\_blank">18626576</a>). Negatively regulates the pathway by phosphorylating the scaffold protein TAXBP1 and thus promoting the assembly of the A20/TNFAIP3 ubiquitin-editing complex (composed of A20/TNFAIP3, TAX1BP1, and the E3 ligases ITCH and RNF11) (PubMed:<a href="http://www.uniprot.org/citations/21765415" target="\_blank">21765415</a>). Therefore, CHUK plays a key role in the negative feedback of NF-kappa-B canonical signaling to limit inflammatory gene activation. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes (PubMed:<a href="http://www.uniprot.org/citations/20501937" target="\_blank">20501937</a>). In turn, these complexes regulate genes encoding molecules involved in B-cell survival and lymphoid organogenesis. Participates also in the negative feedback of the non-canonical NF- kappa-B signaling pathway by phosphorylating and destabilizing MAP3K14/NIK. Within the nucleus, phosphorylates CREBBP and consequently increases both its transcriptional and histone acetyltransferase activities (PubMed:<a href="http://www.uniprot.org/citations/17434128" target="\_blank">17434128</a>). Modulates chromatin accessibility at NF- kappa-B-responsive promoters by phosphorylating histones H3 at 'Ser-10' that are subsequently acetylated at 'Lys-14' by CREBBP (PubMed:<a href="http://www.uniprot.org/citations/12789342" target="\_blank">12789342</a>). Additionally, phosphorylates the CREBBP-interacting protein NCOA3. Also phosphorylates FOXO3 and may regulate this pro- apoptotic transcription factor (PubMed:<a href="http://www.uniprot.org/citations/15084260" target="\_blank">15084260</a>). Phosphorylates RIPK1 at 'Ser-25' which represses its kinase activity and consequently prevents TNF-mediated RIPK1-dependent cell death (By similarity). Phosphorylates AMBRA1 following mitophagy induction, promoting AMBRA1 interaction with ATG8 family proteins and its mitophagic activity (PubMed:<a href="http://www.uniprot.org/citations/30217973" target="\_blank">30217973</a>).

#### **Cellular Location**

Cytoplasm. Nucleus Note=Shuttles between the cytoplasm and the nucleus

#### **Tissue Location**

Widely expressed.

### **Phospho-CHUK(T23) Antibody Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**Phospho-CHUK(T23) Antibody Blocking peptide - Images****Phospho-CHUK(T23) Antibody Blocking peptide - Background**

CHUK is a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquination pathway, thereby activating the transcription factor.

**Phospho-CHUK(T23) Antibody Blocking peptide - References**

Park, K.J., et al., Mol. Cell 18(1):71-82 (2005).Qing, G., et al., J. Biol. Chem. 280(11):9765-9768 (2005).Xiao, G., et al., J. Biol. Chem. 279(29):30099-30105 (2004).Gu, L., et al., J. Biol. Chem. 279(50):52141-52149 (2004).Buss, H., et al., J. Biol. Chem. 279(53):55633-55643 (2004).