

Phospho-APS(Y629) Antibody Blocking peptide
Synthetic peptide
Catalog # BP3703a

Specification

Phospho-APS(Y629) Antibody Blocking peptide - Product Information

Primary Accession [O14492](#)

Phospho-APS(Y629) Antibody Blocking peptide - Additional Information

Gene ID 10603

Other Names

SH2B adapter protein 2, Adapter protein with pleckstrin homology and Src homology 2 domains, SH2 and PH domain-containing adapter protein APS, SH2B2, APS

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Phospho-APS(Y629) Antibody Blocking peptide - Protein Information

Name SH2B2

Synonyms APS

Function

Adapter protein for several members of the tyrosine kinase receptor family. Involved in multiple signaling pathways. May be involved in coupling from immunoreceptor to Ras signaling. Acts as a negative regulator of cytokine signaling in collaboration with CBL. Binds to EPOR and suppresses EPO-induced STAT5 activation, possibly through a masking effect on STAT5 docking sites in EPOR. Suppresses PDGF-induced mitogenesis. May induce cytoskeletal reorganization via interaction with VAV3.

Cellular Location

Cytoplasm. Cell membrane. Note=Cytoplasmic before PDGF stimulation. After PDGF stimulation, localized at the cell membrane and peripheral region

Tissue Location

Expressed in spleen, prostate, testis, uterus, small intestine and skeletal muscle. Among hematopoietic cell lines, expressed exclusively in B-cells. Not expressed in most tumor cell lines.

Phospho-APS(Y629) Antibody Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Phospho-APS(Y629) Antibody Blocking peptide - Images**Phospho-APS(Y629) Antibody Blocking peptide - Background**

APS is expressed in B lymphocytes and contains pleckstrin homology and src homology 2 (SH2) domains. In Burkitt lymphoma cell lines, it is tyrosine phosphorylated in response to B cell receptor stimulation. Because it binds Shc independent of stimulation and Grb2 after stimulation, it appears to play a role in signal transduction from the receptor to Shc/Grb2.

Phospho-APS(Y629) Antibody Blocking peptide - References

Li,M., Endocrinology 148 (4), 1615-1621 (2007)Katsanakis,K.D., J. Biol. Chem. 280 (45), 37827-37832 (2005)Iseki,M., Biochem. Biophys. Res. Commun. 330 (3), 1005-1013 (2005)