

**Phospho-rat TSC2(T1373) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP3833a****Specification**

---

**Phospho-rat TSC2(T1373) Blocking Peptide - Product Information**

Primary Accession [P49816](#)  
Other Accession [NP\\_036812.2](#)

**Phospho-rat TSC2(T1373) Blocking Peptide - Additional Information**

**Gene ID** 24855

**Other Names**

Tuberin, Tuberous sclerosis 2 protein homolog, Tsc2

**Target/Specificity**

The synthetic peptide sequence is selected from aa 1366-1380 of RAT Tsc2

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Phospho-rat TSC2(T1373) Blocking Peptide - Protein Information**

**Name** Tsc2 {ECO:0000303|PubMed:8519695, ECO:0000312|RGD:3908}

**Function**

Catalytic component of the TSC-TBC complex, a multiprotein complex that acts as a negative regulator of the canonical mTORC1 complex, an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:<a href="http://www.uniprot.org/citations/12172553" target="\_blank">12172553</a>). Within the TSC-TBC complex, TSC2 acts as a GTPase-activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1 (By similarity). In absence of nutrients, the TSC-TBC complex inhibits mTORC1, thereby preventing phosphorylation of ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) by the mTORC1 signaling (By similarity). The TSC-TBC complex is inactivated in response to nutrients, relieving inhibition of mTORC1 (By similarity). Involved in microtubule-mediated protein transport via its ability to regulate mTORC1 signaling (PubMed:<a href="http://www.uniprot.org/citations/16707451" target="\_blank">16707451</a>). Also stimulates the intrinsic GTPase activity of the Ras-related proteins RAP1A and RAB5 (PubMed:<a href="http://www.uniprot.org/citations/9045618" target="\_blank">9045618</a>).

**Cellular Location**

Lysosome membrane {ECO:0000250|UniProtKB:P49815}; Peripheral membrane protein {ECO:0000250|UniProtKB:P49815}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:P49815}.  
Note=Recruited to lysosomal membranes in a RHEB-dependent process in absence of nutrients (By similarity). In response to insulin signaling and phosphorylation by PKB/AKT1, the complex dissociates from lysosomal membranes and relocalizes to the cytosol (By similarity) {ECO:0000250|UniProtKB:P49815}

**Tissue Location**

CNS, uterus, heart, skeletal muscle, kidney and spleen.

**Phospho-rat TSC2(T1373) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**Phospho-rat TSC2(T1373) Blocking Peptide - Images****Phospho-rat TSC2(T1373) Blocking Peptide - Background**

acts as a tumor suppressor; may play a role in cell cycle regulation; acute phase response, and negative regulation of cell proliferation [RGD].

**Phospho-rat TSC2(T1373) Blocking Peptide - References**

Larson, Y., et al. J. Biol. Chem. 285(32):24987-24998(2010)  
Sajankila, S.P., et al. Mol. Cell. Biochem. 338 (1-2), 233-239 (2010) :  
Inoue, H., et al. Biosci. Biotechnol. Biochem. 73(11):2488-2493(2009)  
Di Nardo, A., et al. J. Neurosci. 29(18):5926-5937(2009)  
Shiono, M., et al. Oncogene 27(52):6690-6697(2008)