

GNAT2 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP4873b**Specification**

GNAT2 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession [P19087](#)

GNAT2 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 2780

Other Names

Guanine nucleotide-binding protein G(t) subunit alpha-2, Transducin alpha-2 chain, GNAT2, GNATC

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

GNAT2 Antibody (C-term) Blocking Peptide - Protein Information

Name GNAT2

Synonyms GNATC

Function

Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. Transducin is an amplifier and one of the transducers of a visual impulse that performs the coupling between rhodopsin and cGMP- phosphodiesterase.

Cellular Location

Cell projection, cilium, photoreceptor outer segment {ECO:0000250|UniProtKB:P50149}. Photoreceptor inner segment {ECO:0000250|UniProtKB:P50149}. Note=Localizes mainly in the outer segment in the dark-adapted state, whereas is translocated to the inner part of the photoreceptors in the light-adapted state. During dark- adapted conditions, in the presence of UNC119 mislocalizes from the outer segment to the inner part of rod photoreceptors which leads to decreased photoreceptor damage caused by light {ECO:0000250|UniProtKB:P50149}

Tissue Location

Retinal rod outer segment.

GNAT2 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

GNAT2 Antibody (C-term) Blocking Peptide - Images

GNAT2 Antibody (C-term) Blocking Peptide - Background

Transducin is a 3-subunit guanine nucleotide-binding protein (G protein) which stimulates the coupling of rhodopsin and cGMP-phosphodiesterase during visual impulses. The transducin alpha subunits in rods and cones are encoded by separate genes. This gene encodes the alpha subunit in cones.

GNAT2 Antibody (C-term) Blocking Peptide - References

Thiadens, A.A., et al. Ophthalmology 116(10):1984-1989(2009) Luttrell, L.M. Mol. Biotechnol. 39(3):239-264(2008) Oldham, W.M., et al. Nat. Struct. Mol. Biol. 13(9):772-777(2006) Rosenberg, T., et al. Invest. Ophthalmol. Vis. Sci. 45(12):4256-4262(2004)