

## VLDLR Antibody (Center) Blocking peptide

Synthetic peptide Catalog # BP5754c

### **Specification**

## VLDLR Antibody (Center) Blocking peptide - Product Information

Primary Accession P98155
Other Accession NP\_003374.3

## VLDLR Antibody (Center) Blocking peptide - Additional Information

**Gene ID 7436** 

### **Other Names**

Very low-density lipoprotein receptor, VLDL receptor, VLDL-R, VLDLR

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### VLDLR Antibody (Center) Blocking peptide - Protein Information

#### Name VLDLR

## **Function**

Multifunctional cell surface receptor that binds VLDL and transports it into cells by endocytosis and therefore plays an important role in energy metabolism. Binds also to a wide range of other molecules including Reelin/RELN or apolipoprotein E/APOE- containing ligands as well as clusterin/CLU (PubMed:<a href="http://www.uniprot.org/citations/24381170" target="\_blank">24381170</a>, PubMed:<a href="http://www.uniprot.org/citations/30873003" target="\_blank">30873003</a>). In the off-state of the pathway, forms homooligomers or heterooligomers with LRP8 (PubMed:<a href="http://www.uniprot.org/citations/30873003" target="\_blank">30873003</a>). Upon binding to ligands, homooligomers are rearranged to higher order receptor clusters that transmit the extracellular RELN signal to intracellular signaling processes by binding to DAB1 (PubMed:<a href="http://www.uniprot.org/citations/30873003" target="\_blank">30873003</a>). This interaction results in phosphorylation of DAB1 leading to the ultimate cell responses required for the correct positioning of newly generated neurons. Later, mediates a stop signal for migrating neurons, preventing them from entering the marginal zone (By similarity).

#### **Cellular Location**

Cell membrane; Single-pass type I membrane protein Membrane, clathrin-coated pit; Single-pass



type I membrane protein

### **Tissue Location**

Abundant in heart and skeletal muscle; also ovary and kidney; not in liver

### **VLDLR Antibody (Center) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

VLDLR Antibody (Center) Blocking peptide - Images

## VLDLR Antibody (Center) Blocking peptide - Background

The low density lipoprotein receptor (LDLR) gene familyconsists of cell surface proteins involved in receptor-mediatedendocytosis of specific ligands. This gene encodes a lipoprotein receptor that is a member of the LDLR family and plays important roles in VLDL-triglyceride metabolism and the reelin signaling pathway. Mutations in this gene cause VLDLR-associated cerebellar hypoplasia. Alternative splicing generates multiple transcript variants encoding distinct isoforms for this gene. [provided by Ref Seq].

# **VLDLR Antibody (Center) Blocking peptide - References**

Sakai, K., et al. Brain Res. 1276, 11-21 (2009)Francis, P.J., et al. J. Med. Genet. 46(5):300-307(2009)Ananyeva, N.M., et al. Blood Coagul. Fibrinolysis 19(6):543-555(2008)Turkmen, S., et al. Eur. J. Hum. Genet. 16(9):1070-1074(2008)