

ADCY10 Antibody (Center) Blocking peptide

Synthetic peptide Catalog # BP5862c

Specification

ADCY10 Antibody (Center) Blocking peptide - Product Information

Primary Accession Q96PN6

Other Accession <u>NP_001161221.1</u>, <u>NP_060887.2</u>

ADCY10 Antibody (Center) Blocking peptide - Additional Information

Gene ID 55811

Other Names

Adenylate cyclase type 10, AH-related protein, Adenylate cyclase homolog, Germ cell soluble adenylyl cyclase, hsAC, sAC, Testicular soluble adenylyl cyclase, ADCY10, SAC

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ADCY10 Antibody (Center) Blocking peptide - Protein Information

Name ADCY10

Synonyms SAC

Function

Catalyzes the formation of the signaling molecule cAMP (PubMed:12609998, PubMed:15659711, PubMed:24616449, PubMed:25040695, PubMed:25040695, PubMed:24567411). May function as sensor that mediates responses to changes in cellular bicarbonate and CO(2) levels (PubMed:15659711/a>, PubMed:17591988). Has a critical role in mammalian spermatogenesis by producing the cAMP which regulates cAMP-responsive nuclear factors indispensable for sperm maturation in the epididymis. Induces capacitation, the maturational process that sperm undergo prior to fertilization (By similarity). Involved in ciliary beat regulation (PubMed:17591988/a>).



Tel: 858.875.1900 Fax: 858.875.1999

Cellular Location

Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cytoskeleton. Cytoplasm, perinuclear region. Nucleus. Cell projection, cilium. Cytoplasm Mitochondrion. Note=Distributed to subcellular compartments containing cAMP targets. Found as a plasma membrane-associated protein, protein concentrated in the perinuclear region and protein colocalized with actin or tubulin

Tissue Location

Detected in airway epithelial cells and testis (at protein level) (PubMed:17591988). Weakly expressed in multiple tissues Expressed in brain, heart, kidney, liver, lung, pancreas, peripheral blood leukocytes, placenta, skeletal muscle, stomach, thymus, airway epithelial cells, duodenum, jejunum and ileum. Very low level of expression in bone.

ADCY10 Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

ADCY10 Antibody (Center) Blocking peptide - Images