

# ARF1 Antibody (Center) Blocking peptide

Synthetic peptide Catalog # BP5871c

### **Specification**

## ARF1 Antibody (Center) Blocking peptide - Product Information

Primary Accession <u>P84077</u>

Other Accession NP\_001019398.1, NP\_001649.1, NP\_001019399.1, NP\_001019397.1

### ARF1 Antibody (Center) Blocking peptide - Additional Information

Gene ID 375

**Other Names** 

ADP-ribosylation factor 1, ARF1

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**


This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## ARF1 Antibody (Center) Blocking peptide - Protein Information

#### Name ARF1

# **Function**

Small GTPase involved in protein trafficking between different compartments (PubMed:<a href="http://www.uniprot.org/citations/8253837" target="\_blank">8253837</a>). Modulates vesicle budding and uncoating within the Golgi complex (PubMed:<a href="http://www.uniprot.org/citations/8253837" target="\_blank">8253837</a>). In its GTP-bound form, triggers the recruitment of coatomer proteins to the Golgi membrane (PubMed:<a href="http://www.uniprot.org/citations/8253837" target="\_blank">8253837</a>). The hydrolysis of ARF1-bound GTP, which is mediated by ARFGAPs proteins, is required for dissociation of coat proteins from Golgi membranes and vesicles (PubMed:<a href="http://www.uniprot.org/citations/8253837" target="\_blank">8253837</a>). The GTP- bound form interacts with PICK1 to limit PICK1-mediated inhibition of Arp2/3 complex activity; the function is linked to AMPA receptor (AMPAR) trafficking, regulation of synaptic plasticity of excitatory synapses and spine shrinkage during long-term depression (LTD) (By similarity). Plays a key role in the regulation of intestinal stem cells and gut microbiota, and is essential for maintaining intestinal homeostasis (By similarity). Plays also a critical role in mast cell expansion but not in mast cell maturation by facilitating optimal mTORC1 activation (By similarity).





Tel: 858.875.1900 Fax: 858.875.1999

### **Cellular Location**

Golgi apparatus membrane; Lipid-anchor; Cytoplasmic side. Synapse, synaptosome {ECO:0000250|UniProtKB:P84079}. Postsynaptic density {ECO:0000250|UniProtKB:P84079}. Note=In the GDP-bound form, associates transiently with the membranes via its myristoylated N-terminus where guanine nucleotide-exchange factor (GEF)-mediated nucleotide exchange occurs (By similarity). Following nucleotide exchange, the GTP-bound form undergoes a conformational change, leading to the exposure of a myristoylated N-terminal amphipathic helix that provides stable membrane anchorage (By similarity). {ECO:0000250|UniProtKB:P84080}

### ARF1 Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

ARF1 Antibody (Center) Blocking peptide - Images