

P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide
Synthetic peptide
Catalog # BP6111a**Specification**

P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide - Product InformationPrimary Accession [P08183](#)**P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide - Additional Information****Gene ID** 5243**Other Names**

Multidrug resistance protein 1, ATP-binding cassette sub-family B member 1, P-glycoprotein 1, CD243, ABCB1, MDR1, PGY1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6111a](/product/products/AP6111a) was selected from the Center region of human ABCB1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide - Protein Information**Name** ABCB1 ([HGNC:40](#))**Synonyms** MDR1, PGY1**Function**

Translocates drugs and phospholipids across the membrane (PubMed: [8898203](http://www.uniprot.org/citations/8898203), PubMed: [2897240](http://www.uniprot.org/citations/2897240), PubMed: [9038218](http://www.uniprot.org/citations/9038218), PubMed: [35970996](http://www.uniprot.org/citations/35970996)). Catalyzes the flop of phospholipids from the cytoplasmic to the exoplasmic leaflet of the apical membrane. Participates mainly to the flop of phosphatidylcholine, phosphatidylethanolamine, beta-D-glucosylceramides and sphingomyelins (PubMed: [8898203](http://www.uniprot.org/citations/8898203)).

Energy-dependent efflux pump responsible for decreased drug accumulation in multidrug-resistant cells (PubMed:2897240, PubMed:9038218, PubMed:35970996).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000255|PROSITE-ProRule:PRU00441} Apical cell membrane. Cytoplasm Note=ABCB1 localization is influenced by C1orf115 expression levels (plasma membrane versus cytoplasm). Localized to the apical membrane of enterocytes (PubMed:28408210).

Tissue Location

Expressed in small intestine (PubMed:28408210). Expressed in liver, kidney and brain.

P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide - Images

P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide - Background

The membrane-associated ABCB1 protein is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance. ABCB1 is an ATP-dependent drug efflux pump for xenobiotic compounds with broad substrate specificity. It is responsible for decreased drug accumulation in multidrug-resistant cells and often mediates the development of resistance to anticancer drugs. This protein also functions as a transporter in the blood-brain barrier.

P-Glycoprotein (ABCB1) Antibody (Center) Blocking peptide - References

Saito, S., et al., J. Hum. Genet. 47(1):38-50 (2002).Kerb, R., et al., Pharmacogenomics 2(1):51-64 (2001).Cascorbi, I., et al., Clin. Pharmacol. Ther. 69(3):169-174 (2001).Hoffmeyer, S., et al., Proc. Natl. Acad. Sci. U.S.A. 97(7):3473-3478 (2000).Mickley, L.A., et al., Blood 91(5):1749-1756 (1998).