

**NRG2 Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP6223b****Specification**

---

**NRG2 Antibody (Center) Blocking Peptide - Product Information**Primary Accession [O14511](#)**NRG2 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 9542**Other Names**

Pro-neuregulin-2, membrane-bound isoform, Pro-NRG2, Neuregulin-2, NRG-2, Divergent of neuregulin-1, DON-1, Neural- and thymus-derived activator for ERBB kinases, NTAK, NRG2, NTAK

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6223b](/product/products/AP6223b) was selected from the Center region of human NRG2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**NRG2 Antibody (Center) Blocking Peptide - Protein Information****Name** NRG2**Synonyms** NTAK**Function**

Direct ligand for ERBB3 and ERBB4 tyrosine kinase receptors. Concomitantly recruits ERBB1 and ERBB2 coreceptors, resulting in ligand-stimulated tyrosine phosphorylation and activation of the ERBB receptors. May also promote the heterodimerization with the EGF receptor.

**Cellular Location**

[Pro-neuregulin-2, membrane-bound isoform]: Cell membrane; Single-pass type I membrane protein. Note=Does not seem to be active.

**Tissue Location**

Restricted to the cerebellum in the adult.

### **NRG2 Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **NRG2 Antibody (Center) Blocking Peptide - Images**

### **NRG2 Antibody (Center) Blocking Peptide - Background**

Neuregulin 2 (NRG2) is a novel member of the neuregulin family of growth and differentiation factors. Through interaction with the ErbB family of receptors, NRG2 induces the growth and differentiation of epithelial, neuronal, glial, and other types of cells. The gene consists of 12 exons and the genomic structure is similar to that of neuregulin 1 (NRG1), another member of the neuregulin family of ligands. NRG1 and NRG2 mediate distinct biological processes by acting at different sites in tissues and eliciting different biological responses in cells. There are six NRG2 alternatively spliced transcripts which are variable between the epidermal growth factor-like (EGF-like) and the transmembrane domains. The gene is located close to the region for demyelinating Charcot-Marie-Tooth disease locus, but is not responsible for this disease.