

**DSCR3 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP6319b****Specification**

---

**DSCR3 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O14972](#)**DSCR3 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 10311

**Other Names**

Down syndrome critical region protein 3, Down syndrome critical region protein A, DSCR3, DCRA, DSCRA

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6319b](/product/products/AP6319b) was selected from the C-term region of human DSCR3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**DSCR3 Antibody (C-term) Blocking Peptide - Protein Information**Name VPS26C ([HGNC:3044](#))**Function**

Acts as a component of the retriever complex. The retriever complex is a heterotrimeric complex related to retromer cargo-selective complex (CSC) and essential for retromer-independent retrieval and recycling of numerous cargos such as integrin alpha-5/beta-1 (ITGA5:ITGB1) (PubMed: [28892079](http://www.uniprot.org/citations/28892079)). The recruitment of the retriever complex to the endosomal membrane involves CCC and WASH complexes (PubMed: [28892079](http://www.uniprot.org/citations/28892079)). In the endosomes, drives the retriever and recycling of NxxY-motif-containing cargo proteins by coupling to SNX17, a cargo essential for the homeostatic maintenance of numerous cell surface proteins associated with processes that include cell migration, cell adhesion, nutrient supply and cell signaling (PubMed: [28892079](http://www.uniprot.org/citations/28892079)).

**Cellular Location**

Endosome.

**Tissue Location**

Ubiquitously expressed.

**DSCR3 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**DSCR3 Antibody (C-term) Blocking Peptide - Images****DSCR3 Antibody (C-term) Blocking Peptide - Background**

The region of chromosome 21 between genes CBR and ERG (CBR-ERG region), which spans 2.5 Mb on 21q22.2, has been defined by analysis of patients with partial trisomy 21. It contributes significantly to the pathogenesis of many characteristics of Down syndrome, including morphological features, hypotonia, and mental retardation. The DSCR3 (Down syndrome critical region gene 3) gene is found in this region and is predicted to contain eight exons. DSCR3 is expressed in most tissues examined.

**DSCR3 Antibody (C-term) Blocking Peptide - References**

Dahmane, N., et al., Genomics 48(1):12-23 (1998). Nakamura, A., et al., J. Biochem. 122(4):872-877 (1997).