

CLCA1 Antibody (Center K598) Blocking Peptide
Synthetic peptide
Catalog # BP6335e**Specification**

CLCA1 Antibody (Center K598) Blocking Peptide - Product Information

Primary Accession [A8K7I4](#)
Other Accession [Q9UPC6](#)

CLCA1 Antibody (Center K598) Blocking Peptide - Additional Information

Gene ID 1179

Other Names

Calcium-activated chloride channel regulator 1, 34--, Calcium-activated chloride channel family member 1, hCLCA1, Calcium-activated chloride channel protein 1, CaCC-1, hCaCC-1, CLCA1, CACC1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6335e](/product/products/AP6335e) was selected from the Center region of human CLCA1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CLCA1 Antibody (Center K598) Blocking Peptide - Protein Information

Name CLCA1

Synonyms CACC1

Function

May be involved in mediating calcium-activated chloride conductance. May play critical roles in goblet cell metaplasia, mucus hypersecretion, cystic fibrosis and AHR. May be involved in the regulation of mucus production and/or secretion by goblet cells. Involved in the regulation of tissue inflammation in the innate immune response. May play a role as a tumor suppressor. Induces MUC5AC.

Cellular Location

Secreted, extracellular space. Cell membrane; Peripheral membrane protein; Extracellular side.
Note=Protein that remains attached to the plasma membrane appeared to be predominantly localized to microvilli

Tissue Location

Highly expressed in small intestine and colon namely in intestinal basal crypt epithelia and goblet cells, and appendix. Weakly expressed in uterus, testis and kidney. Expressed in the airways epithelium of both asthmatic and healthy patients Expressed in the bronchial epithelium, especially in mucus-producing goblet cells. Expressed in normal turbinate mucosa and nasal polyp Expressed in.

CLCA1 Antibody (Center K598) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CLCA1 Antibody (Center K598) Blocking Peptide - Images**CLCA1 Antibody (Center K598) Blocking Peptide - Background**

CLCA1 belongs to the calcium sensitive chloride conductance protein family. To date, all members of this gene family map to the same site on chromosome 1p31-p22 and share high degrees of homology in size, sequence and predicted structure, but differ significantly in their tissue distributions. This gene product is expressed predominantly in the small intestine and colon mucosa, and may be involved in secretory or absorptive processes.

CLCA1 Antibody (Center K598) Blocking Peptide - References

Ritzka, M., et al., Hum. Genet. 115(6):483-491 (2004).Kamada, F., et al., Genes Immun. 5(7):540-547 (2004).Greenwood, I.A., et al., J. Biol. Chem. 277(25):22119-22122 (2002).Loewen, M.E., et al., Biochem. Biophys. Res. Commun. 298(4):531-536 (2002).Toda, M., et al., J. Allergy Clin. Immunol. 109(2):246-250 (2002).