

RAD9 Antibody (C-term P334) Blocking Peptide
Synthetic peptide
Catalog # BP6357c**Specification**

RAD9 Antibody (C-term P334) Blocking Peptide - Product InformationPrimary Accession [Q99638](#)**RAD9 Antibody (C-term P334) Blocking Peptide - Additional Information****Gene ID** 5883**Other Names**

Cell cycle checkpoint control protein RAD9A, hRAD9, DNA repair exonuclease rad9 homolog A, RAD9A

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6357c](/products/AP6357c) was selected from the C-term region of human Rad9. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

RAD9 Antibody (C-term P334) Blocking Peptide - Protein Information**Name** RAD9A**Function**

Component of the 9-1-1 cell-cycle checkpoint response complex that plays a major role in DNA repair (PubMed: [10713044](http://www.uniprot.org/citations/10713044), PubMed: [17575048](http://www.uniprot.org/citations/17575048), PubMed: [20545769](http://www.uniprot.org/citations/20545769), PubMed: [21659603](http://www.uniprot.org/citations/21659603), PubMed: [31135337](http://www.uniprot.org/citations/31135337)). The 9-1-1 complex is recruited to DNA lesion upon damage by the RAD17- replication factor C (RFC) clamp loader complex (PubMed: [21659603](http://www.uniprot.org/citations/21659603)). Acts then as a sliding clamp platform on DNA for several proteins involved in long-patch base excision repair (LP-BER) (PubMed: [21659603](http://www.uniprot.org/citations/21659603))

target="_blank">21659603). The 9-1-1 complex stimulates DNA polymerase beta (POLB) activity by increasing its affinity for the 3'-OH end of the primer-template and stabilizes POLB to those sites where LP-BER proceeds; endonuclease FEN1 cleavage activity on substrates with double, nick, or gap flaps of distinct sequences and lengths; and DNA ligase I (LIG1) on long-patch base excision repair substrates (PubMed:21659603). The 9-1-1 complex is necessary for the recruitment of RHNO1 to sites of double-stranded breaks (DSB) occurring during the S phase (PubMed:21659603). RAD9A possesses 3'->5' double stranded DNA exonuclease activity (PubMed:10713044).

Cellular Location

Nucleus.

RAD9 Antibody (C-term P334) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

RAD9 Antibody (C-term P334) Blocking Peptide - Images

RAD9 Antibody (C-term P334) Blocking Peptide - Background

Rad9 is highly similar to *Schizosaccharomyces pombe* rad9, a cell cycle checkpoint protein required for cell cycle arrest and DNA damage repair in response to DNA damage. This protein is found to possess 3' to 5' exonuclease activity, which may contribute to its role in sensing and repairing DNA damage. It forms a checkpoint protein complex with RAD1 and HUS1. This complex is recruited by checkpoint protein RAD17 to the sites of DNA damage, which is thought to be important for triggering the checkpoint-signaling cascade.

RAD9 Antibody (C-term P334) Blocking Peptide - References

Maniwa, Y., et al., *Cancer* 103(1):126-132 (2005). Wang, W., et al., *Proc. Natl. Acad. Sci. U.S.A.* 101(48):16762-16767 (2004). Lindsey-Boltz, L.A., et al., (er) *Nucleic Acids Res.* 32(15):4524-4530 (2004). Toueille, M., et al., (er) *Nucleic Acids Res.* 32(11):3316-3324 (2004). Loegering, D., et al., *J. Biol. Chem.* 279(18):18641-18647 (2004).