

TRAP Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP6545a**Specification****TRAP Antibody (N-term) Blocking Peptide - Product Information**

Primary Accession [P29965](#)

TRAP Antibody (N-term) Blocking Peptide - Additional Information**Gene ID 959****Other Names**

CD40 ligand, CD40-L, T-cell antigen Gp39, TNF-related activation protein, TRAP, Tumor necrosis factor ligand superfamily member 5, CD154, CD40 ligand, membrane form, CD40 ligand, soluble form, CD40LG, CD40L, TNFSF5, TRAP

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6545a](#) was selected from the N-term region of human TRAP. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TRAP Antibody (N-term) Blocking Peptide - Protein Information

Name CD40LG

Synonyms CD40L, TNFSF5, TRAP

Function

Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed:[1280226](http://www.uniprot.org/citations/1280226), PubMed:[31331973](http://www.uniprot.org/citations/31331973)). Costimulates T-cell proliferation and cytokine production (PubMed:[8617933](http://www.uniprot.org/citations/8617933)). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:[8617933](http://www.uniprot.org/citations/8617933)). Induces the

activation of NF-kappa-B (PubMed:15067037, PubMed:31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching (By similarity).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Cell surface

Tissue Location

Specifically expressed on activated CD4+ T- lymphocytes

TRAP Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

TRAP Antibody (N-term) Blocking Peptide - Images**TRAP Antibody (N-term) Blocking Peptide - Background**

TRAP is expressed on the surface of T cells. It regulates B cell function by engaging CD40 on the B cell surface. A defect in its gene results in an inability to undergo immunoglobulin class switch and is associated with hyper-IgM syndrome.

TRAP Antibody (N-term) Blocking Peptide - References

Volmar,C.H., Exp. Cell Res. 315 (13), 2265-2274 (2009)Chai,H., Surgery 146 (1), 5-11 (2009)