

AGO1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP6558a**Specification****AGO1 Antibody (N-term) Blocking Peptide - Product Information**

Primary Accession [Q9UL18](#)

AGO1 Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 26523

Other Names

Protein argonaute-1, Argonaute1, hAgo1, Argonaute RISC catalytic component 1, Eukaryotic translation initiation factor 2C 1, eIF-2C 1, eIF2C 1, Putative RNA-binding protein Q99, AGO1, EIF2C1

Target/Specificity

The synthetic peptide sequence used to generate the antibody <[/products/AP6558a](#)>AP6558a was selected from the N-term region of human AGO1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

AGO1 Antibody (N-term) Blocking Peptide - Protein Information

Name AGO1

Synonyms EIF2C1

Function

Required for RNA-mediated gene silencing (RNAi). Binds to short RNAs such as microRNAs (miRNAs) or short interfering RNAs (siRNAs), and represses the translation of mRNAs which are complementary to them. Lacks endonuclease activity and does not appear to cleave target mRNAs. Also required for transcriptional gene silencing (TGS) of promoter regions which are complementary to bound short antogene RNAs (agRNAs).

Cellular Location

Cytoplasm, P-body

AGO1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

AGO1 Antibody (N-term) Blocking Peptide - Images

AGO1 Antibody (N-term) Blocking Peptide - Background

AGO1 is a member of the Argonaute family of proteins which play a role in RNA interference. The protein is highly basic, and contains a PAZ domain and a PIWI domain. It may interact with dicer1 and play a role in short-interfering-RNA-mediated gene silencing.

AGO1 Antibody (N-term) Blocking Peptide - References

Weinmann,L., Cell 136 (3), 496-507 (2009) Horikawa,Y., Clin. Cancer Res. 14 (23), 7956-7962 (2008)