

CDCA3 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP6614a**Specification**

CDCA3 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q99618](#)**CDCA3 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 83461**Other Names**

Cell division cycle-associated protein 3, Gene-rich cluster protein C8, Trigger of mitotic entry protein 1, TOME-1, CDCA3, C8, GRCC8, TOME1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6614a](/products/AP6614a) was selected from the N-term region of human CDCA3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CDCA3 Antibody (N-term) Blocking Peptide - Protein Information**Name** CDCA3**Synonyms** C8, GRCC8, TOME1**Function**

F-box-like protein which is required for entry into mitosis. Acts by participating in E3 ligase complexes that mediate the ubiquitination and degradation of WEE1 kinase at G2/M phase (By similarity).

Cellular Location

Cytoplasm, cytosol.

CDCA3 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CDCA3 Antibody (N-term) Blocking Peptide - Images**CDCA3 Antibody (N-term) Blocking Peptide - Background**

CDCA3 is a F-box-like protein which is required for entry into mitosis. It acts by participating to E3 ligase complexes that mediate the ubiquitination and degradation of WEE1 kinase at G2/M phase.

CDCA3 Antibody (N-term) Blocking Peptide - References

Daub H., Mol. Cell 31:438-448(2008)