

## CRTC3 Antibody (C-term) Blocking Peptide Synthetic peptide

Catalog # BP6633b

## Specification

# **CRTC3 Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession

<u>Q6UUV7</u>

## CRTC3 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 64784

**Other Names** 

CREB-regulated transcription coactivator 3, Transducer of regulated cAMP response element-binding protein 3, TORC-3, Transducer of CREB protein 3, CRTC3, TORC3

## Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP6633b>AP6633b</a> was selected from the C-term region of human CRTC3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **CRTC3 Antibody (C-term) Blocking Peptide - Protein Information**

Name CRTC3

Synonyms TORC3

## Function

Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. Potent coactivator of PPARGC1A and inducer of mitochondrial biogenesis in muscle cells. Also coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR).

**Cellular Location** 



Nucleus. Cytoplasm. Note=Appears to be mainly nuclear (PubMed:15454081). Translocates to the nucleus following adenylyl cyclase or MAP kinase activation (PubMed:30611118)

## **Tissue Location**

Predominantly expressed in B and T lymphocytes. Highest levels in lung. Also expressed in brain, colon, heart, kidney, ovary, and prostate. Weak expression in liver, pancreas, muscle, small intestine, spleen and stomach.

## **CRTC3 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

### <u>Blocking Peptides</u>

### CRTC3 Antibody (C-term) Blocking Peptide - Images

## **CRTC3 Antibody (C-term) Blocking Peptide - Background**

CRTC3 is a transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. The protein acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. The protein nnhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. It is a potent coactivator of PPARGC1A and inducer of mitochondrial biogenesis in muscle cells. It ia also a coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR).

### **CRTC3 Antibody (C-term) Blocking Peptide - References**

Fehr,A., Genes Chromosomes Cancer 47 (3), 203-206 (2008)Wu,Z., Proc. Natl. Acad. Sci. U.S.A. 103 (39), 14379-14384 (2006)