

DREV Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP6636b

Specification

DREV Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

Q9H1A3

DREV Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 51108

Other Names

Methyltransferase-like protein 9, DORA reverse strand protein, DREV, DREV1, METTL9, DREV

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP6636b was selected from the C-term region of human DREV. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DREV Antibody (C-term) Blocking Peptide - Protein Information

Name METTL9 {ECO:0000303|PubMed:33563959, ECO:0000312|HGNC:HGNC:24586}

Function

Protein-histidine N-methyltransferase that specifically catalyzes 1-methylhistidine (pros-methylhistidine) methylation of target proteins (PubMed:33563959, PubMed:34562450). Mediates methylation of proteins with a His-x-His (HxH) motif (where 'x' is preferably a small amino acid) (PubMed:33563959). Catalyzes methylation of target proteins such as S100A9, NDUFB3, SLC39A5, SLC39A7, ARMC6 and DNAJB12; 1-methylhistidine modification may affect the binding of zinc and other metals to its target proteins (PubMed:3356395934562450). Constitutes the main methyltransferase for the 1-methylhistidine modification in cell (PubMed:<a href="http://www.uniprot.org/citations/33563959"



Tel: 858.875.1900 Fax: 858.875.1999

target="_blank">33563959).

Cellular Location

Endoplasmic reticulum Mitochondrion. Note=Colocalizes with membranous compartments such as the endoplasmic reticulum and mitochondria.

DREV Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

DREV Antibody (C-term) Blocking Peptide - Images

DREV Antibody (C-term) Blocking Peptide - Background

DREV belongs to the DREV family.

DREV Antibody (C-term) Blocking Peptide - References

Shu, K.X., Colloids Surf B Biointerfaces 52 (1), 22-30 (2006)Tsang, H.T., Genomics 88 (3), 333-346 (2006)