

**KPNA3 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP6767b**

### Specification

#### **KPNA3 Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession [O00505](#)

#### **KPNA3 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 3839

##### **Other Names**

Importin subunit alpha-4, Importin alpha Q2, Qip2, Karyopherin subunit alpha-3, SRP1-gamma, KPNA3, QIP2

##### **Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6767b](#) was selected from the C-term region of human KPNA3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

##### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

##### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

##### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

#### **KPNA3 Antibody (C-term) Blocking Peptide - Protein Information**

**Name** KPNA3

**Synonyms** QIP2

##### **Function**

Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1. Binds specifically and directly to substrates containing either a simple or bipartite NLS motif. Docking of the importin/substrate complex to the nuclear pore complex (NPC) is mediated by KPNB1 through binding to nucleoporin FxFG repeats and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components separate and importin-alpha and -beta are re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran from importin. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. In vitro, mediates the

nuclear import of human cytomegalovirus UL84 by recognizing a non-classical NLS. Recognizes NLSs of influenza A virus nucleoprotein probably through ARM repeats 7-9.

**Cellular Location**

Cytoplasm. Nucleus

**Tissue Location**

Ubiquitous. Highest levels in heart and skeletal muscle

**KPNA3 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**KPNA3 Antibody (C-term) Blocking Peptide - Images****KPNA3 Antibody (C-term) Blocking Peptide - Background**

KPNA3 belongs to the importin alpha family, and is involved in nuclear protein import.

**KPNA3 Antibody (C-term) Blocking Peptide - References**

Singh, A.P., et.al., Cell 131 (3), 492-504 (2007)