

FGA Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP6847a**Specification**

FGA Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P02671](#)**FGA Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 2243**Other Names**

Fibrinogen alpha chain, Fibrinopeptide A, Fibrinogen alpha chain, FGA

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6847a](/products/AP6847a) was selected from the N-term region of human FGA. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

FGA Antibody (N-term) Blocking Peptide - Protein Information**Name** FGA**Function**

Cleaved by the protease thrombin to yield monomers which, together with fibrinogen beta (FGB) and fibrinogen gamma (FGG), polymerize to form an insoluble fibrin matrix. Fibrin has a major function in hemostasis as one of the primary components of blood clots. In addition, functions during the early stages of wound repair to stabilize the lesion and guide cell migration during re-epithelialization. Was originally thought to be essential for platelet aggregation, based on in vitro studies using anticoagulated blood. However, subsequent studies have shown that it is not absolutely required for thrombus formation in vivo. Enhances expression of SELP in activated platelets via an ITGB3-dependent pathway. Maternal fibrinogen is essential for successful pregnancy. Fibrin deposition is also associated with infection, where it protects against IFNG-mediated hemorrhage. May also facilitate the immune response via both innate and T-cell mediated pathways.

Cellular Location

Secreted

Tissue Location

Detected in blood plasma (at protein level).

FGA Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

FGA Antibody (N-term) Blocking Peptide - Images**FGA Antibody (N-term) Blocking Peptide - Background**

FGA is the alpha component of fibrinogen, a blood-borne glycoprotein comprised of three pairs of nonidentical polypeptide chains. Following vascular injury, fibrinogen is cleaved by thrombin to form fibrin which is the most abundant component of blood clots. In addition, various cleavage products of fibrinogen and fibrin regulate cell adhesion and spreading, display vasoconstrictor and chemotactic activities, and are mitogens for several cell types.

FGA Antibody (N-term) Blocking Peptide - References

Thomas,V.A.,et.al., Biochemistry 37 (8), 2291-2298 (1998)