

## EPB41L4B Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP6851b

## **Specification**

# EPB41L4B Antibody (C-term) Blocking Peptide - Product Information

**Primary Accession** 

**Q9H329** 

# EPB41L4B Antibody (C-term) Blocking Peptide - Additional Information

**Gene ID 54566** 

#### **Other Names**

Band 41-like protein 4B, FERM-containing protein CG1, Protein EHM2, EPB41L4B, EHM2

### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP6851b>AP6851b</a> was selected from the C-term region of human EPB41L4B. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## EPB41L4B Antibody (C-term) Blocking Peptide - Protein Information

Name EPB41L4B (<u>HGNC:19818</u>)

### **Function**

Up-regulates the activity of the Rho guanine nucleotide exchange factor ARHGEF18 (By similarity). Involved in the regulation of the circumferential actomyosin belt in epithelial cells (PubMed:<a href="http://www.uniprot.org/citations/22006950" target="\_blank">22006950</a>). Promotes cellular adhesion, migration and motility in vitro and may play a role in wound healing (PubMed:<a href="http://www.uniprot.org/citations/23664528" target="\_blank">23664528</a>). May have a role in mediating cytoskeletal changes associated with steroid- induced cell differentiation (PubMed:<a href="http://www.uniprot.org/citations/14521927" target="\_blank">14521927</a>).

## **Cellular Location**

Cytoplasm. Cell junction, tight junction Note=Accumulates along apical cell-cell boundaries and is also detected in the cytoplasm in a punctate manner.



Tel: 858.875.1900 Fax: 858.875.1999

## **Tissue Location**

Expressed at higher levels in acute wounds than chronic wounds with increased expression in healing wounds, especially at the leading wound edge (PubMed:23664528). Isoform 1 is highly expressed in brain. Isoform 2 is highly expressed in testis with lower levels in prostate and breast (PubMed:14521927)

# EPB41L4B Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

EPB41L4B Antibody (C-term) Blocking Peptide - Images

EPB41L4B Antibody (C-term) Blocking Peptide - References

Leyne, M., et.al., Unpublished