

**GTF2I Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP6881b****Specification**

---

**GTF2I Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [P78347](#)**GTF2I Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 2969**Other Names**

General transcription factor II-I, GTFII-I, TFII-I, Bruton tyrosine kinase-associated protein 135, BAP-135, BTK-associated protein 135, SRF-Phox1-interacting protein, SPIN, Williams-Beuren syndrome chromosomal region 6 protein, GTF2I, BAP135, WBSCR6

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP6881b](/products/AP6881b) was selected from the C-term region of human GTF2I. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**GTF2I Antibody (C-term) Blocking Peptide - Protein Information****Name** GTF2I**Synonyms** BAP135, WBSCR6**Function**

Interacts with the basal transcription machinery by coordinating the formation of a multiprotein complex at the C-FOS promoter, and linking specific signal responsive activator complexes. Promotes the formation of stable high-order complexes of SRF and PHOX1 and interacts cooperatively with PHOX1 to promote serum-inducible transcription of a reporter gene driven by the C-FOS serum response element (SRE). Acts as a coregulator for USF1 by binding independently two promoter elements, a pyrimidine-rich initiator (Inr) and an upstream E-box. Required for the formation of functional ARID3A DNA-binding complexes and for activation of immunoglobulin heavy-chain transcription upon B-lymphocyte activation.



**Cellular Location**

Cytoplasm. Nucleus {ECO:0000255|PROSITE-ProRule:PRU00484, ECO:0000269|PubMed:10373551} Note=Colocalizes with BTK in the cytoplasm

**Tissue Location**

Ubiquitous. Isoform 1 is strongly expressed in fetal brain, weakly in adult brain, muscle, and lymphoblasts and is almost undetectable in other adult tissues, while the other isoforms are equally expressed in all adult tissues

**GTF2I Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**GTF2I Antibody (C-term) Blocking Peptide - Images****GTF2I Antibody (C-term) Blocking Peptide - Background**

GTF2I is a multifunctional phosphoprotein with roles in transcription and signal transduction. It is deleted in Williams-Beuren syndrome, a multisystem developmental disorder caused by the deletion of contiguous genes at chromosome 7q11.23.

**GTF2I Antibody (C-term) Blocking Peptide - References**

Roy,A.L.,et.al., EMBO J. 16 (23), 7091-7104 (1997)