

HADHA Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP6882b**Specification**

HADHA Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [P40939](#)**HADHA Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 3030**Other Names**

Trifunctional enzyme subunit alpha, mitochondrial, 78 kDa gastrin-binding protein, TP-alpha, Long-chain enoyl-CoA hydratase, Long chain 3-hydroxyacyl-CoA dehydrogenase, HADHA, HADH

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6882b](/products/AP6882b) was selected from the C-term region of human HADHA. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

HADHA Antibody (C-term) Blocking Peptide - Protein Information**Name** HADHA**Synonyms** HADH**Function**

Mitochondrial trifunctional enzyme catalyzes the last three of the four reactions of the mitochondrial beta-oxidation pathway (PubMed: [8135828](http://www.uniprot.org/citations/8135828), PubMed: [1550553](http://www.uniprot.org/citations/1550553), PubMed: [29915090](http://www.uniprot.org/citations/29915090), PubMed: [30850536](http://www.uniprot.org/citations/30850536)). The mitochondrial beta-oxidation pathway is the major energy-producing process in tissues and is performed through four consecutive reactions breaking down fatty acids into acetyl-CoA (PubMed: [29915090](http://www.uniprot.org/citations/29915090)).

Among the enzymes involved in this pathway, the trifunctional enzyme exhibits specificity for long-chain fatty acids (PubMed:30850536). Mitochondrial trifunctional enzyme is a heterotetrameric complex composed of two proteins, the trifunctional enzyme subunit alpha/HADHA described here carries the 2,3-enoyl-CoA hydratase and the 3-hydroxyacyl-CoA dehydrogenase activities while the trifunctional enzyme subunit beta/HADHB bears the 3-ketoacyl-CoA thiolase activity (PubMed:8135828, PubMed:29915090, PubMed:30850536). Independently of the subunit beta, the trifunctional enzyme subunit alpha/HADHA also has a monolysocardiolipin acyltransferase activity (PubMed:23152787). It acylates monolysocardiolipin into cardiolipin, a major mitochondrial membrane phospholipid which plays a key role in apoptosis and supports mitochondrial respiratory chain complexes in the generation of ATP (PubMed:23152787). Allows the acylation of monolysocardiolipin with different acyl-CoA substrates including oleoyl-CoA for which it displays the highest activity (PubMed:23152787).

Cellular Location

Mitochondrion. Mitochondrion inner membrane Note=Protein stability and association with mitochondrion inner membrane do not require HADHB.

HADHA Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

HADHA Antibody (C-term) Blocking Peptide - Images

HADHA Antibody (C-term) Blocking Peptide - Background

HADHA is the alpha subunit of the mitochondrial trifunctional protein, which catalyzes the last three steps of mitochondrial beta-oxidation of long chain fatty acids. The mitochondrial membrane-bound heterocomplex is composed of four alpha and four beta subunits, with the alpha subunit catalyzing the 3-hydroxyacyl-CoA dehydrogenase and enoyl-CoA hydratase activities.

HADHA Antibody (C-term) Blocking Peptide - References

Sims,H.F., et.al., Proc. Natl. Acad. Sci. U.S.A. 92 (3), 841-845 (1995)