

## **HGD Antibody (C-term) Blocking Peptide**

Synthetic peptide Catalog # BP6883b

## **Specification**

## **HGD Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession

**Q93099** 

# **HGD Antibody (C-term) Blocking Peptide - Additional Information**

**Gene ID 3081** 

#### **Other Names**

Homogentisate 1, 2-dioxygenase, Homogentisate oxygenase, Homogentisic acid oxidase, Homogentisicase, HGD, HGO

# **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a href=/products/AP6883b>AP6883b</a> was selected from the C-term region of human HGD. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **HGD Antibody (C-term) Blocking Peptide - Protein Information**

Name HGD

Synonyms HGO

### **Function**

Catalyzes the conversion of homogentisate to maleylacetoacetate.

#### **Tissue Location**

Highest expression in the prostate, small intestine, colon, kidney and liver

### **HGD Antibody (C-term) Blocking Peptide - Protocols**





Provided below are standard protocols that you may find useful for product applications.

# • Blocking Peptides

**HGD Antibody (C-term) Blocking Peptide - Images** 

**HGD Antibody (C-term) Blocking Peptide - Background** 

Homogentisate 1,2-dioxygenase (HGD) gene mutations are the molecular cause of alkaptonuria, a rare hereditary disorder of the phenylalanine catabolism.

**HGD Antibody (C-term) Blocking Peptide - References** 

Abdulrazzag, Y.M., et.al., Ann. Hum. Genet. 73 (1), 125-130 (2009)