

FMO3 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP6901a**Specification**

FMO3 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P31513](#)**FMO3 Antibody (N-term) Blocking Peptide - Additional Information**

Gene ID 2328

Other Names

Dimethylaniline monooxygenase [N-oxide-forming] 3, Dimethylaniline oxidase 3, FMO II, FMO form 2, Hepatic flavin-containing monooxygenase 3, FMO 3, Trimethylamine monooxygenase, FMO3

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP6901a](/products/AP6901a) was selected from the N-term region of human FMO3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

FMO3 Antibody (N-term) Blocking Peptide - Protein Information**Name** FMO3**Function**

Essential hepatic enzyme that catalyzes the oxygenation of a wide variety of nitrogen- and sulfur-containing compounds including drugs as well as dietary compounds (PubMed: [10759686](http://www.uniprot.org/citations/10759686), PubMed: [30381441](http://www.uniprot.org/citations/30381441), PubMed: [32156684](http://www.uniprot.org/citations/32156684)). Plays an important role in the metabolism of trimethylamine (TMA), via the production of trimethylamine N-oxide (TMAO) metabolite (PubMed: [9776311](http://www.uniprot.org/citations/9776311)). TMA is generated by the action of gut microbiota using dietary precursors such as choline, choline containing compounds, betaine or L-carnitine. By regulating TMAO concentration, FMO3 directly impacts both platelet responsiveness and rate of thrombus formation (PubMed: [29981269](http://www.uniprot.org/citations/29981269))

target="_blank">29981269).

Cellular Location

Microsome membrane {ECO:0000250|UniProtKB:P32417}; Single-pass membrane protein.
Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P32417}; Single-pass membrane protein

Tissue Location

Liver.

FMO3 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

FMO3 Antibody (N-term) Blocking Peptide - Images**FMO3 Antibody (N-term) Blocking Peptide - Background**

FMO3 is involved in the oxidative metabolism of a variety of xenobiotics such as drugs and pesticides. It N-oxygenates primary aliphatic alkylamines as well as secondary and tertiary amines. It plays an important role in the metabolism of trimethylamine (TMA), via the production of TMA N-oxide (TMAO). Is also able to perform S-oxidation when acting on sulfide compounds.

FMO3 Antibody (N-term) Blocking Peptide - References

Allerston,C.K., et.al., Mol. Genet. Metab. 98 (1-2), 198-202 (2009)