

APBB1IP Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP6977c

Specification

APBB1IP Antibody (Center) Blocking Peptide - Product Information

Primary Accession

Q7Z5R6

APBB1IP Antibody (Center) Blocking Peptide - Additional Information

Gene ID 54518

Other Names

Amyloid beta A4 precursor protein-binding family B member 1-interacting protein, APBB1-interacting protein 1, Proline-rich EVH1 ligand 1, PREL-1, Proline-rich protein 73, Rap1-GTP-interacting adapter molecule, RIAM, Retinoic acid-responsive proline-rich protein 1, RARP-1, APBB1IP, PREL1, RARP1, RIAM

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP6977c was selected from the Center region of human APBB1IP. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

APBB1IP Antibody (Center) Blocking Peptide - Protein Information

Name APBB1IP

Synonyms PREL1, RARP1, RIAM

Function

Appears to function in the signal transduction from Ras activation to actin cytoskeletal remodeling. Suppresses insulin-induced promoter activities through AP1 and SRE. Mediates Rap1-induced adhesion.

Cellular Location

Cell membrane; Peripheral membrane protein. Cell projection, lamellipodium Cell junction, focal adhesion. Cytoplasm, cytoskeleton. Note=Colocalizes with ENA/VASP proteins at lamellipodia tips



Tel: 858.875.1900 Fax: 858.875.1999

and focal adhesions, and F-actin at the leading edge. At the membrane surface, associates, via the PH domain, preferentially with the inositol phosphates, PtdIns(5)P and PtdIns(3)P. This binding appears to be necessary for the efficient interaction of the RA domain to Ras-GTPases (By similarity).

Tissue Location

Widely expressed with high expression in thymus, spleen, lymph node, bone marrow and peripheral leukocytes

APBB1IP Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

APBB1IP Antibody (Center) Blocking Peptide - Images

APBB1IP Antibody (Center) Blocking Peptide - Background

APBB1IP appears to function in the signal transduction from Ras activation to actin cytoskeletal remodeling, suppresses insulin-induced promoter activities through AP1 and SRE and mediates Rap1-induced adhesion.

APBB1IP Antibody (Center) Blocking Peptide - References

Morgan, A.R., et.al., Am. J. Med. Genet. B Neuropsychiatr. Genet. 144B (6), 762-770(2007)