

CHK Antibody (N-term) Blocking Peptide

Synthetic peptide Catalog # BP7064a

Specification

CHK Antibody (N-term) Blocking Peptide - Product Information

Primary Accession

P35790

CHK Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 1119

Other Names

Choline kinase alpha, CK, CHETK-alpha, Ethanolamine kinase, EK, CHKA, CHK, CKI

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP7064a was selected from the N-term region of human CHK. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CHK Antibody (N-term) Blocking Peptide - Protein Information

Name CHKA

Synonyms CHK, CKI {ECO:0000303|PubMed:1618328}

Function

Plays a key role in phospholipid biosynthesis by catalyzing the phosphorylation of free choline to phosphocholine, the first step in phosphatidylcholine biosynthesis (PubMed:19915674, PubMed:34077757, PubMed:17007874). Also phosphorylates ethanolamine, thereby contributing to phosphatidylethanolamine biosynthesis (PubMed:19915674, PubMed:17007874). Has higher activity with choline (PubMed:19915674, PubMed:17007874



target="_blank">17007874). May contribute to tumor cell growth (PubMed:19915674).

Cellular Location Cytoplasm, cytosol.

CHK Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

CHK Antibody (N-term) Blocking Peptide - Images

CHK Antibody (N-term) Blocking Peptide - Background

The dominant pathway for biosynthesis of phosphatidylcholine occurs through the CDP-choline pathway. Choline kinase alpha (CHK) is the first enzyme in the pathway and may thereby play an upstream regulatory role in lipid transport and metabolism. CHK additionally catalyzes phosphorylation of ethanolamine.