

CCRK Antibody (C-term H288) Blocking Peptide
Synthetic peptide
Catalog # BP7119b**Specification**

CCRK Antibody (C-term H288) Blocking Peptide - Product InformationPrimary Accession [Q8IZL9](#)**CCRK Antibody (C-term H288) Blocking Peptide - Additional Information****Gene ID** 23552**Other Names**

Cyclin-dependent kinase 20, CDK-activating kinase p42, CAK-kinase p42, Cell cycle-related kinase, Cell division protein kinase 20, Cyclin-dependent protein kinase H, Cyclin-kinase-activating kinase p42, CDK20, CCRK, CDCH

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7119b](/products/AP7119b) was selected from the C-term region of human CCRK. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CCRK Antibody (C-term H288) Blocking Peptide - Protein Information**Name** CDK20**Synonyms** CCRK, CDCH**Function**

Required for high-level Shh responses in the developing neural tube. Together with TBC1D32, controls the structure of the primary cilium by coordinating assembly of the ciliary membrane and axoneme, allowing GLI2 to be properly activated in response to SHH signaling (By similarity). Involved in cell growth. Activates CDK2, a kinase involved in the control of the cell cycle, by phosphorylating residue 'Thr-160'.

Cellular Location

Nucleus. Cytoplasm. Cell projection, cilium

CCRK Antibody (C-term H288) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CCRK Antibody (C-term H288) Blocking Peptide - Images

CCRK Antibody (C-term H288) Blocking Peptide - Background

CCRK contains a kinase domain most closely related to the cyclin-dependent protein kinases. This protein is involved in cell growth and activates CDK2, a kinase involved in the control of the cell cycle, by phosphorylating residue Thr-160.

CCRK Antibody (C-term H288) Blocking Peptide - References

Fujii, H., et al., Biochem. Biophys. Res. Commun. 322(3):1052-1058 (2004). Liu, Y., et al., J. Biol. Chem. 279(6):4507-4514 (2004).