

IGHM Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP7470b**Specification**

IGHM Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [P01871](#)**IGHM Antibody (C-term) Blocking Peptide - Additional Information****Other Names**

Ig mu chain C region, IGHM

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

IGHM Antibody (C-term) Blocking Peptide - Protein Information**Name** IGHM {ECO:0000303|PubMed:11340299, ECO:0000303|Ref.14}**Function**

Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:22158414, PubMed:20176268). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268).

Cellular Location

[Isoform 1]: Secreted. Note=During differentiation, B-lymphocytes switch from expression of membrane-bound IgM to secretion of IgM.

IGHM Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

IGHM Antibody (C-term) Blocking Peptide - Images

IGHM Antibody (C-term) Blocking Peptide - References

Dorai H., Gillies S.D. Nucleic Acids Res. 17:6412-6412(1989) Rabbitts T.H. Nucleic Acids Res. 9:4509-4524(1981) Watanabe S. Hoppe-Seyler's Z. Physiol. Chem. 354:1505-1509(1973)