

**PKM1 Antibody (C-term L398) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP7476b**

**Specification**

**PKM1 Antibody (C-term L398) Blocking peptide - Product Information**

Primary Accession [P14618](#)

**PKM1 Antibody (C-term L398) Blocking peptide - Additional Information**

**Gene ID** 5315

**Other Names**

Pyruvate kinase PKM, Cytosolic thyroid hormone-binding protein, CTHBP, Opa-interacting protein 3, OIP-3, Pyruvate kinase 2/3, Pyruvate kinase muscle isozyme, Thyroid hormone-binding protein 1, THBP1, Tumor M2-PK, p58, PKM, OIP3, PK2, PK3, PKM2

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP7476b was selected from the C-term region of human PKM2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**PKM1 Antibody (C-term L398) Blocking peptide - Protein Information**

**Name** PKM

**Synonyms** OIP3 {ECO:0000303|PubMed:9466265}, PK2,

**Function**

Catalyzes the final rate-limiting step of glycolysis by mediating the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP (PubMed:<a href="http://www.uniprot.org/citations/15996096" target="\_blank">15996096</a>, PubMed:<a href="http://www.uniprot.org/citations/1854723" target="\_blank">1854723</a>, PubMed:<a href="http://www.uniprot.org/citations/20847263" target="\_blank">20847263</a>). The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production (PubMed:<a href="http://www.uniprot.org/citations/15996096" target="\_blank">15996096</a>, PubMed:<a href="http://www.uniprot.org/citations/1854723" target="\_blank">1854723</a>,

PubMed:<a href="http://www.uniprot.org/citations/20847263" target="\_blank">20847263</a>). The transition between the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and survival (PubMed:<a href="http://www.uniprot.org/citations/15996096" target="\_blank">15996096</a>, PubMed:<a href="http://www.uniprot.org/citations/1854723" target="\_blank">1854723</a>, PubMed:<a href="http://www.uniprot.org/citations/20847263" target="\_blank">20847263</a>).

### **Cellular Location**

[Isoform M2]: Cytoplasm. Nucleus Note=Translocates to the nucleus in response to various signals, such as EGF receptor activation or apoptotic stimuli (PubMed:17308100, PubMed:22056988, PubMed:24120661). Nuclear translocation is promoted by acetylation by EP300 (PubMed:24120661). Deacetylation by SIRT6 promotes its nuclear export in a process dependent of XPO4, thereby suppressing its ability to activate transcription and promote tumorigenesis (PubMed:26787900).

### **Tissue Location**

[Isoform M2]: Specifically expressed in proliferating cells, such as embryonic stem cells, embryonic carcinoma cells, as well as cancer cells.

## **PKM1 Antibody (C-term L398) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

## **PKM1 Antibody (C-term L398) Blocking peptide - Images**

## **PKM1 Antibody (C-term L398) Blocking peptide - Background**

PKM1 is a pyruvate kinase that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. The protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis.

## **PKM1 Antibody (C-term L398) Blocking peptide - References**

Dombrauckas J.D. *Biochemistry* 44:9417-9429(2005) Daub H., Olsen J.V. *Mol. Cell* 31:438-448(2008)  
Ashizawa K., McPhie P. *Biochemistry* 30:7105-7111(1991)