

ITK Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP7711c**Specification**

ITK Antibody (Center) Blocking Peptide - Product Information

Primary Accession
Other Accession

[Q08881](#)
[ITK_HUMAN](#)

ITK Antibody (Center) Blocking Peptide - Additional Information

Gene ID 3702

Other Names

Tyrosine-protein kinase ITK/TSK, Interleukin-2-inducible T-cell kinase, IL-2-inducible T-cell kinase, Kinase EMT, T-cell-specific kinase, Tyrosine-protein kinase Lyk, ITK, EMT, LYK

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP7711c](/product/products/AP7711c) was selected from the Center region of human ITK. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ITK Antibody (Center) Blocking Peptide - Protein Information

Name ITK

Synonyms EMT, LYK

Function

Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation lead to the recruitment of ITK to the cell membrane, in the vicinity of the stimulated TCR receptor, where it is phosphorylated by LCK. Phosphorylation leads to ITK autophosphorylation and full activation. Once activated, phosphorylates PLCG1, leading to the activation of this lipase and subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT)

translocates into the nucleus to perform its transcriptional duty. Phosphorylates 2 essential adapter proteins: the linker for activation of T-cells/LAT protein and LCP2. Then, a large number of signaling molecules such as VAV1 are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation (PubMed:12186560, PubMed:12682224, PubMed:21725281). Required for TCR-mediated calcium response in gamma-delta T-cells, may also be involved in the modulation of the transcriptomic signature in the Vgamma2-positive subset of immature gamma-delta T-cells (By similarity). Phosphorylates TBX21 at 'Tyr-530' and mediates its interaction with GATA3 (By similarity).

Cellular Location

Cytoplasm. Nucleus {ECO:0000250|UniProtKB:Q03526}. Note=Localizes in the vicinity of cell surface receptors in the plasma membrane after receptor stimulation

Tissue Location

T-cell lines and natural killer cell lines.

ITK Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ITK Antibody (Center) Blocking Peptide - Images**ITK Antibody (Center) Blocking Peptide - Background**

ITK is an intracellular tyrosine kinase expressed in T-cells. The protein contains both SH2 and SH3 domains which are often found in intracellular kinases. It is thought to play a role in T-cell proliferation and differentiation.

ITK Antibody (Center) Blocking Peptide - References

Brazin, K.N., et al., Proc. Natl. Acad. Sci. U.S.A. 99(4):1899-1904 (2002).Hawkins, J., et al., Protein Expr. Purif. 22(2):211-219 (2001).Janis, E.M., et al., Genomics 23(1):269-271 (1994).Gibson, S., et al., Blood 82(5):1561-1572 (1993).Tanaka, N., et al., FEBS Lett. 324(1):1-5 (1993).