

**CCK4 (PTK7) Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP7800b****Specification**

---

**CCK4 (PTK7) Antibody (C-term) Blocking peptide - Product Information**Primary Accession [Q13308](#)**CCK4 (PTK7) Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 5754**Other Names**

Inactive tyrosine-protein kinase 7, Colon carcinoma kinase 4, CCK-4, Protein-tyrosine kinase 7, Pseudo tyrosine kinase receptor 7, Tyrosine-protein kinase-like 7, PTK7, CCK4

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP7800b](/product/products/AP7800b) was selected from the C-term region of human CCK4 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**CCK4 (PTK7) Antibody (C-term) Blocking peptide - Protein Information****Name** PTK7**Synonyms** CCK4**Function**

Inactive tyrosine kinase involved in Wnt signaling pathway. Component of both the non-canonical (also known as the Wnt/planar cell polarity signaling) and the canonical Wnt signaling pathway. Functions in cell adhesion, cell migration, cell polarity, proliferation, actin cytoskeleton reorganization and apoptosis. Has a role in embryogenesis, epithelial tissue organization and angiogenesis.

**Cellular Location**

Membrane; Single-pass type I membrane protein. Cell junction. Note=Colocalizes with MMP14 at cell junctions. Also localizes at the leading edge of migrating cells

**Tissue Location**

Highly expressed in lung, liver, pancreas, kidney, placenta and melanocytes. Weakly expressed in thyroid gland, ovary, brain, heart and skeletal muscle. Also expressed in erythroleukemia cells. But not expressed in colon

**CCK4 (PTK7) Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**CCK4 (PTK7) Antibody (C-term) Blocking peptide - Images****CCK4 (PTK7) Antibody (C-term) Blocking peptide - Background**

CCK4 may function as a cell adhesion molecule. Although it belongs to the insuline receptor subfamily of the Tyr protein kinases, it likely lacks the catalytic activity of a tyrosine kinase. It may be connected to the pathophysiology of colon carcinomas and/or may represent a tumor progression marker. This Type I membrane protein is highly expressed in lung, liver, pancreas, kidney, placenta and melanocytes, but weakly expressed in thyroid gland, ovary, brain, heart and skeletal muscle, and not in colon. It is also expressed in erythroleukemia cells.

**CCK4 (PTK7) Antibody (C-term) Blocking peptide - References**

Zhang, H., et al., Nat. Biotechnol. 21(6):660-666 (2003).Park, S.K., et al., J. Biochem. 119(2):235-239 (1996).Mossie, K., et al., Oncogene 11(10):2179-2184 (1995).