

# **ARTS1 Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP7859c

## **Specification**

### **ARTS1 Antibody (Center) Blocking Peptide - Product Information**

Primary Accession <u>Q9NZ08</u>

## ARTS1 Antibody (Center) Blocking Peptide - Additional Information

**Gene ID** 51752

#### **Other Names**

Endoplasmic reticulum aminopeptidase 1, 3411-, ARTS-1, Adipocyte-derived leucine aminopeptidase, A-LAP, Aminopeptidase PILS, Puromycin-insensitive leucyl-specific aminopeptidase, PILS-AP, Type 1 tumor necrosis factor receptor shedding aminopeptidase regulator, ERAP1, APPILS, ARTS1, KIAA0525

### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP7859c>AP7859c</a> was selected from the Center region of human ARTS1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **ARTS1 Antibody (Center) Blocking Peptide - Protein Information**

## Name ERAP1

Synonyms APPILS, ARTS1, KIAA0525

#### **Function**

Aminopeptidase that plays a central role in peptide trimming, a step required for the generation of most HLA class I-binding peptides. Peptide trimming is essential to customize longer precursor peptides to fit them to the correct length required for presentation on MHC class I molecules. Strongly prefers substrates 9-16 residues long. Rapidly degrades 13-mer to a 9-mer and then stops. Preferentially hydrolyzes the residue Leu and peptides with a hydrophobic C-terminus, while it has weak activity toward peptides with charged C-terminus. May play a role in the inactivation of peptide hormones. May be involved in the regulation of blood pressure through the inactivation of



angiotensin II and/or the generation of bradykinin in the kidney.

#### **Cellular Location**

Endoplasmic reticulum membrane; Single-pass type II membrane protein

Tissue Location Ubiquitous.

### **ARTS1 Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

**ARTS1 Antibody (Center) Blocking Peptide - Images** 

### ARTS1 Antibody (Center) Blocking Peptide - Background

Aminopeptidases play a role in the metabolism of several peptides that may be involved in blood pressure and the pathogenesis of essential hypertension. Adipocyte-derived leucine aminopeptidase (ALAP) is a member of the M1 family of zinc metallopeptidases.

## **ARTS1 Antibody (Center) Blocking Peptide - References**

Goto, Y., Biochem. J. 416 (1), 109-116 (2008) Fruci, D., J. Cell. Physiol. 216 (3), 742-749 (2008) Adamik, B., Biochem. Biophys. Res. Commun. 371 (3), 505-509 (2008)