

# AAK1 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP7861b

# **Specification**

# AAK1 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

**Q2M2I8** 

# AAK1 Antibody (C-term) Blocking Peptide - Additional Information

**Gene ID 22848** 

#### **Other Names**

AP2-associated protein kinase 1, Adaptor-associated kinase 1, AAK1, KIAA1048

## Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP7861b>AP7861b</a> was selected from the C-term region of human AAK1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

## **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# AAK1 Antibody (C-term) Blocking Peptide - Protein Information

# Name AAK1

### Synonyms KIAA1048

### **Function**

Regulates clathrin-mediated endocytosis by phosphorylating the AP2M1/mu2 subunit of the adaptor protein complex 2 (AP-2) which ensures high affinity binding of AP-2 to cargo membrane proteins during the initial stages of endocytosis (PubMed:<a

href="http://www.uniprot.org/citations/17494869" target="\_blank">17494869</a>, PubMed:<a href="http://www.uniprot.org/citations/11877457" target="\_blank">11877457</a>, PubMed:<a href="http://www.uniprot.org/citations/11877461" target="\_blank">11877461</a>, PubMed:<a href="http://www.uniprot.org/citations/12952931" target="\_blank">12952931</a>, PubMed:<a href="http://www.uniprot.org/citations/14617351" target="\_blank">14617351</a>, PubMed:<a href="http://www.uniprot.org/citations/14617351" target="\_blank">25653444</a>). Isoform 1 and isoform 2 display similar levels of kinase activity towards AP2M1 (PubMed:<a



href="http://www.uniprot.org/citations/17494869" target="\_blank">17494869</a>). Preferentially, may phosphorylate substrates on threonine residues (PubMed:<a href="http://www.uniprot.org/citations/11877457" target="\_blank">11877457</a>, PubMed:<a href="http://www.uniprot.org/citations/18657069" target="\_blank">18657069</a>). Regulates phosphorylation of other AP-2 subunits as well as AP-2 localization and AP-2-mediated internalization of ligand complexes (PubMed:<a href="http://www.uniprot.org/citations/12952931" target="\_blank">12952931</a>/a>). Phosphorylates NUMB and regulates its cellular localization, promoting NUMB localization to endosomes (PubMed:<a href="http://www.uniprot.org/citations/18657069" target="\_blank">18657069</a>). Binds to and stabilizes the activated form of NOTCH1, increases its localization in endosomes and regulates its transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/21464124" target=" blank">21464124</a>).

#### **Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:F1MH24}; Peripheral membrane protein {ECO:0000250|UniProtKB:F1MH24}. Membrane, clathrin-coated pit. Presynapse {ECO:0000250|UniProtKB:P0C1X8}. Note=Active when found in clathrin- coated pits at the plasma membrane. In neuronal cells, enriched at presynaptic terminals. In non-neuronal cells, enriched at leading edge of migrating cells. {ECO:0000250|UniProtKB:P0C1X8}

# **Tissue Location**

Detected in brain, heart and liver. Isoform 1 is the predominant isoform in brain.

# AAK1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

# • Blocking Peptides

AAK1 Antibody (C-term) Blocking Peptide - Images

### AAK1 Antibody (C-term) Blocking Peptide - Background

Adaptor-related protein complex 2 (AP-2 complexes) functions during receptor-mediated endocytosis to trigger clathrin assembly, interact with membrane-bound receptors, and recruit encodytic accessory factors. AAK1 is a member of the SNF1 subfamily of Ser/Thr protein kinases. The protein interacts with and phosphorylates a subunit of the AP-2 complex, which promotes binding of AP-2 to sorting signals found in membrane-bound receptors and subsequent receptor endocytosis. Its kinase activity is stimulated by clathrin.

# AAK1 Antibody (C-term) Blocking Peptide - References

Henderson, D.M., Mol. Biol. Cell 18 (7), 2698-2706 (2007) Takahashi, T., Cancer Res. 66 (24), 11932-11937 (2006) Schmid, E.M., PLoS Biol. 4 (9), E262 (2006)