

**Protein Phosphatase 1 beta (PPP1CB) Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP8433b**

**Specification**

---

**Protein Phosphatase 1 beta (PPP1CB) Antibody (C-term) Blocking peptide - Product Information**

Primary Accession [P62140](#)

**Protein Phosphatase 1 beta (PPP1CB) Antibody (C-term) Blocking peptide - Additional Information**

**Gene ID** 5500

**Other Names**

Serine/threonine-protein phosphatase PP1-beta catalytic subunit, PP-1B, PPP1CD, PPP1CB

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8433b](/product/products/AP8433b) was selected from the C-term region of human PPP1CB. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Protein Phosphatase 1 beta (PPP1CB) Antibody (C-term) Blocking peptide - Protein Information**

**Name** PPP1CB

**Function**

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase (PP1) is essential for cell division, it participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E. Dephosphorylates the 'Ser-418' residue of FOXP3 in

regulatory T-cells (Treg) from patients with rheumatoid arthritis, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:<a href="http://www.uniprot.org/citations/23396208" target="\_blank">23396208</a>).

**Cellular Location**

Cytoplasm. Nucleus. Nucleus, nucleoplasm. Nucleus, nucleolus. Note=Highly mobile in cells and can be relocalized through interaction with targeting subunits. In the presence of PPP1R8 relocalizes from the nucleus to nuclear speckles.

**Protein Phosphatase 1 beta (PPP1CB) Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**Protein Phosphatase 1 beta (PPP1CB) Antibody (C-term) Blocking peptide - Images****Protein Phosphatase 1 beta (PPP1CB) Antibody (C-term) Blocking peptide - Background**

Protein phosphatase-1 (PP1) is 1 of 4 major serine/threonine-specific protein phosphatases involved in the dephosphorylation of a variety of proteins. These enzymes work in opposition to the protein kinases to control the level of phosphorylation. Protein phosphatase (PP1) is essential for cell division, and it participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis, as well as in regulation of ionic conductances and long-term synaptic plasticity. PP1 has 3 catalytic subunits, designated alpha (PPP1CA), beta (PPP1CB), and gamma (PPP1CC).