

## **UBTF Antibody (N-term) Blocking Peptide**

Synthetic peptide Catalog # BP8572a

## **Specification**

# **UBTF Antibody (N-term) Blocking Peptide - Product Information**

**Primary Accession** 

P17480

## **UBTF Antibody (N-term) Blocking Peptide - Additional Information**

**Gene ID** 7343

#### **Other Names**

Nucleolar transcription factor 1, Autoantigen NOR-90, Upstream-binding factor 1, UBF-1, UBFF, UBF, UBF1

# **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a

href=/products/AP8572a>AP8572a</a> was selected from the N-term region of human UBTF. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **UBTF Antibody (N-term) Blocking Peptide - Protein Information**

Name UBTF

Synonyms UBF, UBF1

### **Function**

Recognizes the ribosomal RNA gene promoter and activates transcription mediated by RNA polymerase I (Pol I) through cooperative interactions with the transcription factor SL1/TIF-IB complex. It binds specifically to the upstream control element and can activate Pol I promoter escape.

## **Cellular Location**

Nucleus, nucleolus {ECO:0000250|UniProtKB:P25976}



Tel: 858.875.1900 Fax: 858.875.1999

# **UBTF Antibody (N-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

**UBTF Antibody (N-term) Blocking Peptide - Images** 

## **UBTF Antibody (N-term) Blocking Peptide - Background**

UBF is a nucleolar phosphoprotein with both DNA binding and transactivation domains. Sequence-specific DNA binding to the core and upstream control elements of the human rRNA promoter is mediated through several HMG boxes.

## **UBTF Antibody (N-term) Blocking Peptide - References**

Hempel, W.M., et.al., Mol. Cell. Biol. 16 (2), 557-563 (1996) Voit, R., et.al., Nucleic Acids Res. 23 (14), 2593-2599 (1995)