

FCGR3B Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP8628c

Specification

FCGR3B Antibody (Center) Blocking Peptide - Product Information

Primary Accession

075015

FCGR3B Antibody (Center) Blocking Peptide - Additional Information

Gene ID 2215

Other Names

Low affinity immunoglobulin gamma Fc region receptor III-B, Fc-gamma RIII-beta, Fc-gamma RIII, Fc-gamma RIIIb, FcRIIIb, FcRIIIb, FcR-10, IgG Fc receptor III-1, CD16b, FCGR3B, CD16B, FCG3, FCGR3, IGFR3

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP8628c was selected from the Center region of human FCGR3B. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

FCGR3B Antibody (Center) Blocking Peptide - Protein Information

Name FCGR3B

Synonyms CD16B, FCG3, FCGR3, IGFR3

Function

Receptor for the Fc region of immunoglobulins gamma. Low affinity receptor. Binds complexed or aggregated IgG and also monomeric IgG. Contrary to III-A, is not capable to mediate antibody-dependent cytotoxicity and phagocytosis. May serve as a trap for immune complexes in the peripheral circulation which does not activate neutrophils.

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor. Secreted Note=Secreted after cleavage



Tissue Location

Expressed specifically by polymorphonuclear leukocytes (neutrophils). Also expressed by stimulated eosinophils

FCGR3B Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

FCGR3B Antibody (Center) Blocking Peptide - Images

FCGR3B Antibody (Center) Blocking Peptide - Background

FCGR3B is a receptor for the Fc region of immunoglobulins gamma. Low affinity receptor. It binds complexed or aggregated IgG and also monomeric IgG. Contrary to III-A, is not capable to mediate antibody-dependent cytotoxicity and phagocytosis. It may serve as a trap for immune complexes in the peripheral circulation which does not activate neutrophils.

FCGR3B Antibody (Center) Blocking Peptide - References

Gessner, J.E., et.al., J. Biol. Chem. 270 (3), 1350-1361 (1995) Ho, Y.P., et.al., J. Clin. Periodontol. (2009)