

HIST1H2BM Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP8780a**Specification**

HIST1H2BM Antibody (N-term) Blocking Peptide - Product Information

Primary Accession [O99879](#)
Other Accession [NP_003512](#)

HIST1H2BM Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 8342

Other Names

Histone H2B type 1-M, Histone H2Be, H2B/e, HIST1H2BM, H2BFE

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8780a](/products/AP8780a) was selected from the N-term region of human HIST1H2BM. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

HIST1H2BM Antibody (N-term) Blocking Peptide - Protein Information

Name H2BC14 ([HGNC:4750](#))

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location

Nucleus. Chromosome.

HIST1H2BM Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

HIST1H2BM Antibody (N-term) Blocking Peptide - Images

HIST1H2BM Antibody (N-term) Blocking Peptide - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures.

HIST1H2BM Antibody (N-term) Blocking Peptide - References

Deng,L., et.al., Virology 277 (2), 278-295 (2000)