

WDR44 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP8818b**Specification**

WDR44 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession [Q5JSH3](#)
Other Accession [NP_061918](#)

WDR44 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 54521

Other Names

WD repeat-containing protein 44, Rabphilin-11, WDR44

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8818b](/products/AP8818b) was selected from the C-term region of human WDR44. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

WDR44 Antibody (C-term) Blocking Peptide - Protein Information

Name WDR44 ([HGNC:30512](#))

Synonyms RPH11

Function

Downstream effector for Rab11 which regulates Rab11 intracellular membrane trafficking functions such as endocytic recycling, intracellular ciliogenesis and protein export (PubMed:[31204173](http://www.uniprot.org/citations/31204173)), PubMed:[32344433](http://www.uniprot.org/citations/32344433)). ATK1-mediated phosphorylation of WDR44 induces binding to Rab11 which activates endocytic recycling of transferrin receptor back to the plasma membrane (PubMed:[31204173](http://www.uniprot.org/citations/31204173)). When bound to Rab11, prevents the formation of the ciliogenic Rab11- Rabin8/RAB3IP-RAB11FIP3 complex, therefore inhibiting preciliary trafficking and ciliogenesis (PubMed:[31204173](http://www.uniprot.org/citations/31204173)).

[31204173](http://www.uniprot.org/citations/31204173)). Participates in neo- synthesized protein export by connecting the endoplasmic reticulum (ER) with the endosomal tubule via direct interactions with the integral ER proteins VAPA or VAPB and the endosomal protein GRAFs (GRAF1/ARHGAP26 or GRAF2/ARHGAP10), which facilitates the transfer of proteins such as E-cadherin, MPP14 and CFTR into a Rab8-Rab10-Rab11-dependent export route (PubMed:[32344433](http://www.uniprot.org/citations/32344433)).

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9R037}. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:Q9R037}. Endosome membrane. Golgi apparatus, trans-Golgi network {ECO:0000250|UniProtKB:Q9R037}. Note=Colocalized with RAB11A, RAB8A, RAB10 and MICAL1 on endosomal tubules (PubMed:32344433). Colocalized with RAB11A along microtubules oriented toward lamellipodia {ECO:0000250|UniProtKB:Q9R037, ECO:0000269|PubMed:32344433}

WDR44 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

WDR44 Antibody (C-term) Blocking Peptide - Images

WDR44 Antibody (C-term) Blocking Peptide - Background

WDR44 is a downstream effector for RAB11. It may be involved in vesicle recycling (By similarity).

WDR44 Antibody (C-term) Blocking Peptide - References

Gevaert,K., et.al., Nat. Biotechnol. 21 (5), 566-569 (2003)