

#### **DNM3 Antibody (Center) Blocking Peptide** Synthetic peptide

Catalog # BP8833c

### Specification

# **DNM3 Antibody (Center) Blocking Peptide - Product Information**

Primary Accession

### <u>Q9UQ16</u>

## DNM3 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 26052

Other Names Dynamin-3, Dynamin, testicular, T-dynamin, DNM3, KIAA0820

### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP8833c>AP8833c</a> was selected from the Center region of human DNM3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## DNM3 Antibody (Center) Blocking Peptide - Protein Information

Name DNM3

Synonyms KIAA0820

Function

Microtubule-associated force-producing protein involved in producing microtubule bundles and able to bind and hydrolyze GTP. Most probably involved in vesicular trafficking processes, in particular endocytosis (By similarity).

**Cellular Location** 

Cytoplasm. Cytoplasm, cytoskeleton. Note=Microtubule-associated.

## DNM3 Antibody (Center) Blocking Peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

#### <u>Blocking Peptides</u>

DNM3 Antibody (Center) Blocking Peptide - Images

### **DNM3 Antibody (Center) Blocking Peptide - Background**

Members of the dynamin family, such as DNM3, possess mechanochemical properties involved in actin-membrane processes, predominantly in membrane budding.

### **DNM3 Antibody (Center) Blocking Peptide - References**

Orth, J.D. et.al., Curr. Opin. Cell Biol. 15 (1), 31-39 (2003)