

**PCCB Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP8843c****Specification**

---

**PCCB Antibody (Center) Blocking Peptide - Product Information**Primary Accession [P05166](#)**PCCB Antibody (Center) Blocking Peptide - Additional Information**

Gene ID 5096

**Other Names**

Propionyl-CoA carboxylase beta chain, mitochondrial, PCCase subunit beta, Propanoyl-CoA:carbon dioxide ligase subunit beta, PCCB

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8843c](/products/AP8843c) was selected from the Center region of human PCCB. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**PCCB Antibody (Center) Blocking Peptide - Protein Information**Name PCCB ([HGNC:8654](#))**Function**

This is one of the 2 subunits of the biotin-dependent propionyl-CoA carboxylase (PCC), a mitochondrial enzyme involved in the catabolism of odd chain fatty acids, branched-chain amino acids isoleucine, threonine, methionine, and valine and other metabolites (PubMed: [15890657](http://www.uniprot.org/citations/15890657), PubMed: [6765947](http://www.uniprot.org/citations/6765947)). Propionyl-CoA carboxylase catalyzes the carboxylation of propionyl-CoA/propanoyl-CoA to D-methylmalonyl-CoA/(S)-methylmalonyl-CoA (PubMed: [15890657](http://www.uniprot.org/citations/15890657), PubMed: [6765947](http://www.uniprot.org/citations/6765947)). Within the holoenzyme, the alpha subunit catalyzes the ATP-dependent carboxylation of the biotin carried by the biotin carboxyl carrier (BCC) domain, while the beta subunit then transfers the carboxyl group from carboxylated biotin to propionyl-CoA

(By similarity). Propionyl-CoA carboxylase also significantly acts on butyryl-CoA/butanoyl-CoA, which is converted to ethylmalonyl-CoA/(2S)-ethylmalonyl-CoA at a much lower rate (PubMed:<a href="http://www.uniprot.org/citations/6765947" target="\_blank">6765947</a>). Other alternative minor substrates include (2E)- butenoyl-CoA/crotonoyl-CoA (By similarity).

**Cellular Location**

Mitochondrion matrix

**PCCB Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**PCCB Antibody (Center) Blocking Peptide - Images****PCCB Antibody (Center) Blocking Peptide - Background**

PCCB is a subunit of the propionyl-CoA carboxylase (PCC) enzyme, which is involved in the catabolism of propionyl-CoA. PCC is a mitochondrial enzyme that probably acts as a dodecamer of six alpha subunits and six beta subunits.

**PCCB Antibody (Center) Blocking Peptide - References**

Yang,X., et.al.,Mol. Genet. Metab. 81 (4), 335-342 (2004)