

CYP51A1 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP8874b**Specification**

CYP51A1 Antibody (C-term) Blocking Peptide - Product InformationPrimary Accession [Q16850](#)**CYP51A1 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 1595**Other Names**

Lanosterol 14-alpha demethylase, LDM, CYPLI, Cytochrome P450 51A1, Cytochrome P450-14DM, Cytochrome P45014DM, Cytochrome P450LI, Sterol 14-alpha demethylase, CYP51A1, CYP51

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8874b](/products/AP8874b) was selected from the C-term region of human CYP51A1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CYP51A1 Antibody (C-term) Blocking Peptide - Protein Information**Name** CYP51A1 ([HGNC:2649](#))**Synonyms** CYP51**Function**

Sterol 14alpha-demethylase that plays a critical role in the cholesterol biosynthesis pathway, being cholesterol the major sterol component in mammalian membranes as well as a precursor for bile acid and steroid hormone synthesis (PubMed: [8619637](http://www.uniprot.org/citations/8619637), PubMed: [9559662](http://www.uniprot.org/citations/9559662), PubMed: [20149798](http://www.uniprot.org/citations/20149798)). Cytochrome P450 monooxygenase that catalyzes the three-step oxidative removal of the 14alpha-methyl group (C-32) of sterols such as lanosterol (lanosta-8,24-dien-3beta-ol) and 24,25-dihydrolanosterol (DHL) in the form of formate, and converts the sterols to

4,4-dimethyl-5 α -cholesta-8,14,24-trien-3 β -ol and 4,4- dimethyl-8,14-cholestadien-3 β -ol, respectively, which are intermediates of cholesterol biosynthesis (PubMed:8619637, PubMed:9559662, PubMed:20149798). Can also demethylate substrates not intrinsic to mammals, such as eburicol (24-methylene-24,25-dihydrolanosterol), but at a lower rate than DHL (PubMed:9559662).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q64654}; Single-pass membrane protein. Microsome membrane {ECO:0000250|UniProtKB:Q64654}; Single-pass membrane protein

Tissue Location

Ubiquitously expressed with highest levels in testis, ovary, adrenal, prostate, liver, kidney and lung

CYP51A1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CYP51A1 Antibody (C-term) Blocking Peptide - Images

CYP51A1 Antibody (C-term) Blocking Peptide - Background

CYP51A1 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum protein participates in the synthesis of cholesterol by catalyzing the removal of the 14 α -methyl group from lanosterol.

CYP51A1 Antibody (C-term) Blocking Peptide - References

Wang,Y., et.al., J. Biol. Chem. 283 (39), 26332-26339 (2008)