

GOT2 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP8968a**Specification**

GOT2 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P00505](#)**GOT2 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 2806**Other Names**

Aspartate aminotransferase, mitochondrial, mAspAT, Fatty acid-binding protein, FABP-1, Glutamate oxaloacetate transaminase 2, Kynurenine aminotransferase 4, Kynurenine aminotransferase IV, Kynurenine--oxoglutarate transaminase 4, Kynurenine--oxoglutarate transaminase IV, Plasma membrane-associated fatty acid-binding protein, FABPpm, Transaminase A, GOT2

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8968a](/products/AP8968a) was selected from the N-term region of human GOT2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

GOT2 Antibody (N-term) Blocking Peptide - Protein Information**Name** GOT2 ([HGNC:4433](#))**Function**

Catalyzes the irreversible transamination of the L-tryptophan metabolite L-kynurenine to form kynurenic acid (KA). As a member of the malate-aspartate shuttle, it has a key role in the intracellular NAD(H) redox balance. Is important for metabolite exchange between mitochondria and cytosol, and for amino acid metabolism. Facilitates cellular uptake of long-chain free fatty acids.

Cellular Location

Mitochondrion matrix. Cell membrane. Note=Exposure to alcohol promotes translocation to the

cell membrane.

GOT2 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

GOT2 Antibody (N-term) Blocking Peptide - Images

GOT2 Antibody (N-term) Blocking Peptide - Background

Glutamic-oxaloacetic transaminase is a pyridoxal phosphate-dependent enzyme which exists in cytoplasmic and inner-membrane mitochondrial forms, GOT1 and GOT2, respectively. GOT plays a role in amino acid metabolism and the urea and tricarboxylic acid cycles. The two enzymes are homodimeric and show close homology.

GOT2 Antibody (N-term) Blocking Peptide - References

Schiele, F., et.al., Clin. Chem. 35 (6), 926-930 (1989) Watazu, Y., et.al., Clin. Chem. 36 (4), 687-689 (1990)