

# **B4GALT3 Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP8977c

# **Specification**

## **B4GALT3 Antibody (Center) Blocking Peptide - Product Information**

**Primary Accession** 

060512

# **B4GALT3 Antibody (Center) Blocking Peptide - Additional Information**

**Gene ID 8703** 

#### **Other Names**

Beta-1, 4-galactosyltransferase 3, Beta-1, 4-GalTase 3, Beta4Gal-T3, b4Gal-T3, 241-, UDP-Gal:beta-GlcNAc beta-1, 4-galactosyltransferase 3, UDP-galactose:beta-N-acetylglucosamine beta-1, 4-galactosyltransferase 3, N-acetyllactosamine synthase, Nal synthase, Beta-N-acetylglucosaminylglycopeptide beta-1, 4-galactosyltransferase, Beta-N-acetylglucosaminyl-glycolipid beta-1, 4-galactosyltransferase, 241-, B4GALT3

### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP8977c>AP8977c</a> was selected from the Center region of human B4GALT3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **B4GALT3 Antibody (Center) Blocking Peptide - Protein Information**

### Name B4GALT3 (HGNC:926)

### **Function**

Responsible for the synthesis of complex-type N-linked oligosaccharides in many glycoproteins as well as the carbohydrate moieties of glycolipids.

#### **Cellular Location**

Golgi apparatus, Golgi stack membrane; Single- pass type II membrane protein. Note=Trans cisternae of Golgi stack

## **Tissue Location**



Found in various tissues. Highest expression in placenta, prostate, testis, ovary, intestine and muscle, and in fetal brain.

# **B4GALT3 Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

**B4GALT3 Antibody (Center) Blocking Peptide - Images** 

# **B4GALT3 Antibody (Center) Blocking Peptide - Background**

B4GALT3 is responsible for the synthesis of complex-type N-linked oligosaccharides in many glycoproteins as well as the carbohydrate moieties of glycolipids.

# **B4GALT3 Antibody (Center) Blocking Peptide - References**

Amado M., et.al., Biochim. Biophys. Acta 1473:35-53(1999).