

**RPA2 Antibody (N-term) ) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP9115a****Specification**

---

**RPA2 Antibody (N-term) ) Blocking Peptide - Product Information**Primary Accession [P15927](#)**RPA2 Antibody (N-term) ) Blocking Peptide - Additional Information****Gene ID** 6118**Other Names**

Replication protein A 32 kDa subunit, RP-A p32, Replication factor A protein 2, RF-A protein 2, Replication protein A 34 kDa subunit, RP-A p34, RPA2, REPA2, RPA32, RPA34

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP9115a](/products/AP9115a) was selected from the N-term region of human RPA2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**RPA2 Antibody (N-term) ) Blocking Peptide - Protein Information****Name** RPA2**Synonyms** REPA2, RPA32, RPA34**Function**

As part of the heterotrimeric replication protein A complex (RPA/RP-A), binds and stabilizes single-stranded DNA intermediates, that form during DNA replication or upon DNA stress. It prevents their reannealing and in parallel, recruits and activates different proteins and complexes involved in DNA metabolism. Thereby, it plays an essential role both in DNA replication and the cellular response to DNA damage. In the cellular response to DNA damage, the RPA complex controls DNA repair and DNA damage checkpoint activation. Through recruitment of ATRIP activates the ATR kinase a master regulator of the DNA damage response. It is required for the recruitment of the DNA double-strand break repair factors RAD51 and RAD52 to chromatin in response to DNA damage. Also recruits to sites of DNA damage proteins like XPA and XPG that are

involved in nucleotide excision repair and is required for this mechanism of DNA repair. Also plays a role in base excision repair (BER) probably through interaction with UNG. Also recruits SMARCAL1/HARP, which is involved in replication fork restart, to sites of DNA damage. May also play a role in telomere maintenance.

**Cellular Location**

Nucleus. Nucleus, PML body. Note=Redistributes to discrete nuclear foci upon DNA damage in an ATR-dependent manner

**RPA2 Antibody (N-term) ) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**RPA2 Antibody (N-term) ) Blocking Peptide - Images****RPA2 Antibody (N-term) ) Blocking Peptide - Background**

RPA2 is required for DNA recombination, repair and replication. The activity of RP-A is mediated by single-stranded DNA binding and protein interactions.

**RPA2 Antibody (N-term) ) Blocking Peptide - References**

Umbricht,C.B., et.al., J. Biol. Chem. 268 (9), 6131-6138 (1993)Oakley,G.G., et.al., Biochemistry 48 (31), 7473-7481 (2009)