

## LOX Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP9127c

### **Specification**

## LOX Antibody (Center) Blocking Peptide - Product Information

Primary Accession

P28300

# LOX Antibody (Center) Blocking Peptide - Additional Information

**Gene ID 4015** 

#### **Other Names**

Protein-lysine 6-oxidase, Lysyl oxidase, LOX

### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/products/AP9127c>AP9127c</a> was selected from the Center region of human LOX. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## LOX Antibody (Center) Blocking Peptide - Protein Information

# **Name LOX**

#### **Function**

Responsible for the post-translational oxidative deamination of peptidyl lysine residues in precursors to fibrous collagen and elastin (PubMed:<a

href="http://www.uniprot.org/citations/26838787" target="\_blank">26838787</a>). Regulator of Ras expression. May play a role in tumor suppression. Plays a role in the aortic wall architecture (By similarity).

#### **Cellular Location**

Secreted. Secreted, extracellular space

### **Tissue Location**

Heart, placenta, skeletal muscle, kidney, lung and pancreas.



# LOX Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

# • Blocking Peptides

LOX Antibody (Center) Blocking Peptide - Images

# LOX Antibody (Center) Blocking Peptide - Background

LOX is an extracellular copper enzyme that initiates the crosslinking of collagens and elastin. The enzyme catalyzes oxidative deamination of the epsilon-amino group in certain lysine and hydroxylysine residues of collagens and lysine residues of elastin. In addition to crosslinking extracellular matrix proteins, the encoded protein may have a role in tumor suppression.

### LOX Antibody (Center) Blocking Peptide - References

Mariani, T.J., et.al., Matrix 12 (3), 242-248 (1992)