

OPRM1 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP9130c**Specification**

OPRM1 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [P35372](#)**OPRM1 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 4988**Other Names**

Mu-type opioid receptor, M-OR-1, MOR-1, Mu opiate receptor, Mu opioid receptor, MOP, hMOP, OPRM1, MOR1

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP9130c](/products/AP9130c) was selected from the Center region of human OPRM1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

OPRM1 Antibody (Center) Blocking Peptide - Protein Information**Name** OPRM1**Synonyms** MOR1**Function**

Receptor for endogenous opioids such as beta-endorphin and endomorphin (PubMed: [10529478](http://www.uniprot.org/citations/10529478), PubMed: [12589820](http://www.uniprot.org/citations/12589820), PubMed: [7891175](http://www.uniprot.org/citations/7891175), PubMed: [7905839](http://www.uniprot.org/citations/7905839), PubMed: [7957926](http://www.uniprot.org/citations/7957926), PubMed: [9689128](http://www.uniprot.org/citations/9689128)). Receptor for natural and synthetic opioids including morphine, heroin, DAMGO, fentanyl, etorphine, buprenorphin and methadone (PubMed: [10529478](http://www.uniprot.org/citations/10529478))

target="_blank">10529478, PubMed:10836142, PubMed:12589820, PubMed:19300905, PubMed:7891175, PubMed:7905839, PubMed:7957926, PubMed:9689128). Also activated by enkephalin peptides, such as Met-enkephalin or Met-enkephalin-Arg-Phe, with higher affinity for Met-enkephalin-Arg-Phe (By similarity). Agonist binding to the receptor induces coupling to an inactive GDP- bound heterotrimeric G-protein complex and subsequent exchange of GDP for GTP in the G-protein alpha subunit leading to dissociation of the G-protein complex with the free GTP-bound G-protein alpha and the G- protein beta-gamma dimer activating downstream cellular effectors (PubMed:7905839). The agonist- and cell type-specific activity is predominantly coupled to pertussis toxin-sensitive G(i) and G(o) G alpha proteins, GNAI1, GNAI2, GNAI3 and GNAO1 isoforms Alpha-1 and Alpha-2, and to a lesser extent to pertussis toxin-insensitive G alpha proteins GNAZ and GNA15 (PubMed:12068084). They mediate an array of downstream cellular responses, including inhibition of adenylate cyclase activity and both N-type and L-type calcium channels, activation of inward rectifying potassium channels, mitogen-activated protein kinase (MAPK), phospholipase C (PLC), phosphoinositide/protein kinase (PKC), phosphoinositide 3-kinase (PI3K) and regulation of NF-kappa-B (By similarity). Also couples to adenylate cyclase stimulatory G alpha proteins (By similarity). The selective temporal coupling to G- proteins and subsequent signaling can be regulated by RGSZ proteins, such as RGS9, RGS17 and RGS4 (By similarity). Phosphorylation by members of the GPRK subfamily of Ser/Thr protein kinases and association with beta-arrestins is involved in short-term receptor desensitization (By similarity). Beta-arrestins associate with the GPRK-phosphorylated receptor and uncouple it from the G-protein thus terminating signal transduction (By similarity). The phosphorylated receptor is internalized through endocytosis via clathrin-coated pits which involves beta-arrestins (By similarity). The activation of the ERK pathway occurs either in a G-protein-dependent or a beta-arrestin- dependent manner and is regulated by agonist-specific receptor phosphorylation (By similarity). Acts as a class A G-protein coupled receptor (GPCR) which dissociates from beta-arrestin at or near the plasma membrane and undergoes rapid recycling (By similarity). Receptor down-regulation pathways are varying with the agonist and occur dependent or independent of G-protein coupling (By similarity). Endogenous ligands induce rapid desensitization, endocytosis and recycling (By similarity). Heterooligomerization with other GPCRs can modulate agonist binding, signaling and trafficking properties (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, axon {ECO:0000250|UniProtKB:P97266}. Perikaryon {ECO:0000250|UniProtKB:P97266}. Cell projection, dendrite {ECO:0000250|UniProtKB:P97266}. Endosome {ECO:0000250|UniProtKB:P97266}. Note=Is rapidly internalized after agonist binding. {ECO:0000250|UniProtKB:P97266}

Tissue Location

Expressed in brain. Isoform 16 and isoform 17 are detected in brain.

OPRM1 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

OPRM1 Antibody (Center) Blocking Peptide - Images

OPRM1 Antibody (Center) Blocking Peptide - Background

OPRM1 is the mu opioid receptor, which is the primary site of action for the most commonly used opioids, including morphine, heroin, fentanyl, and methadone. It is also the primary receptor for endogenous opioid peptides beta-endorphin (see POMC, MIM 176830) and the enkephalins (see PENK, MIM 131330). The OPRM1 receptor is a member of the G protein-coupled receptor family (Bond et al., 1998 [PubMed 9689128]). There are at least 3 types of opioid receptors, mu, kappa (OPRK1; MIM 165196), and delta, each with a distinct pharmacologic profile.

OPRM1 Antibody (Center) Blocking Peptide - References

Bare, L.A., et al., FEBS Lett. 354 (2), 213-216 (1994) Wang, J.B., et al., FEBS Lett. 338 (2), 217-222 (1994)